

Fundamentals in Discrete Convex Analysis

Kazuo Murota¹

Department of Mathematical Informatics
Graduate School of Information Science and Technology
University of Tokyo
113-8656, Tokyo, Japan
`murota@mist.i.u-tokyo.ac.jp`

“Discrete Convex Analysis” is aimed at establishing a novel theoretical framework for solvable discrete optimization problems by means of a combination of the ideas in continuous optimization and combinatorial optimization. The theoretical framework of convex analysis is adapted to discrete settings and the mathematical results in matroid/submodular function theory are generalized. Viewed from the continuous side, the theory can be classified as a theory of convex functions $f : \mathbf{R}^n \rightarrow \mathbf{R}$ that have additional combinatorial properties. Viewed from the discrete side, it is a theory of discrete functions $f : \mathbf{Z}^n \rightarrow \mathbf{Z}$ that enjoy certain nice properties comparable to convexity. Symbolically,

Discrete Convex Analysis = Convex Analysis + Matroid Theory.

The theory puts emphasis on duality and conjugacy as well as on greedy algorithms. This results in a novel duality framework for nonlinear integer programming.

Two convexity concepts, called L-convexity and M-convexity, play primary roles in the present theory, where “L” stands for “Lattice” and “M” for “Matroid.” L-convex functions and M-convex functions are convex functions with additional combinatorial properties distinguished by “L” and “M,” and they are conjugate to each other through a discrete version of the Legendre–Fenchel transformation. L-convex functions and M-convex functions generalize, respectively, the concepts of submodular set functions and base polyhedra of (poly)matroids.

L-convexity and M-convexity prevail in discrete systems.

- In network flow problems, flow and tension are dual objects. Roughly speaking, flow corresponds to M-convexity and tension to L-convexity.
- In matroids, the rank function corresponds to L-convexity and the base family to M-convexity.
- M-matrices correspond to L-convexity, and their inverses to M-convexity. Hence, in a discretization of the Poisson problem of partial differential equations, for example, the differential operator corresponds to L-convexity and the Green function to M-convexity.
- Dirichlet forms in probability theory are essentially the same as quadratic L-convex functions.

This talk is intended to be a brief introduction to the central ideas in discrete convex analysis.

References

1. K. Murota: *Discrete Convex Analysis—An Introduction* (in Japanese), Kyoritsu Publishing Co., Tokyo, 2001.
2. K. Murota: *Discrete Convex Analysis*, SIAM Monographs on Discrete Mathematics and Applications, Vol. 10, SIAM, Philadelphia, 2003.