

# Automating Disease Management Using Answer Set Programming

Zhuo Chen

University of Texas at Dallas, Dallas, Texas, USA  
zxc130130@utdallas.edu

---

## Abstract

Management of chronic diseases such as heart failure, diabetes, and chronic obstructive pulmonary disease (COPD) is a major problem in health care. A standard approach that the medical community has devised to manage widely prevalent chronic diseases such as chronic heart failure (CHF) is to have a committee of experts develop guidelines that all physicians should follow. These guidelines typically consist of a series of complex rules that make recommendations based on a patient's information. Due to their complexity, often the guidelines are either ignored or not complied with at all, which can result in poor medical practices. It is not even clear whether it is humanly possible to follow these guidelines due to their length and complexity. In the case of CHF management, the guidelines run nearly 80 pages. In this paper we describe a physician-advisory system for CHF management that codes the entire set of clinical practice guidelines for CHF using answer set programming. Our approach is based on developing reasoning templates (that we call knowledge patterns) and using these patterns to systemically code the clinical guidelines for CHF as ASP rules. Use of the knowledge patterns greatly facilitates the development of our system. Given a patient's medical information, our system generates a recommendation for treatment just as a human physician would, using the guidelines. Our system will work even in the presence of incomplete information. Our work makes two contributions: (i) it shows that highly complex guidelines can be successfully coded as ASP rules, and (ii) it develops a series of knowledge patterns that facilitate the coding of knowledge expressed in a natural language and that can be used for other application domains.

**1998 ACM Subject Classification** D.1.6 Logic Programming, I.2.3 Logic Programming

**Keywords and phrases** chronic disease management, knowledge pattern, answer set programming, knowledge representation, automated reasoning

**Digital Object Identifier** 10.4230/OASICS.ICLP.2016.22

## 1 Introduction and problem description

Chronic diseases are health conditions that can neither be prevented nor be cured but can only be managed. They have been the major consumer of health-care funds throughout the world. In America alone there are more than 133 million people – which is more than 40% of the U.S. population – who suffer from one or more chronic diseases [17]. In the U.S. they account for 81% of hospital admissions, 91% of prescriptions filled and 76% of all physician visits [1]. Though the list of chronic conditions is long, the top five conditions are: heart disease, cancers, stroke, chronic obstructive pulmonary disease (COPD) and diabetes.

In 2010, 68% of the healthcare spending – more than trillion dollars – went towards the treatment of chronic diseases [5]. The successful management of chronic diseases has two components: (i) self-management by the patients, and (ii) management by physicians while adhering to strict guidelines. The failure of either component will lead to the failure of the whole enterprise for the management of chronic diseases.



© Zhuo Chen;

licensed under Creative Commons License CC-BY

Technical Communications of the 32nd International Conference on Logic Programming (ICLP 2016).

Editors: Manuel Carro, Andy King, Neda Saeedloei, and Marina De Vos; Article No. 22; pp. 22:1–22:10

Open Access Series in Informatics



OASICS Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

In our research, we focus on the second component of CHF management, namely, a Physician Advisory System. This system assists physicians in adhering to the guidelines for managing CHF. The CHF management guidelines are published by the American College of Cardiology Foundation (ACCF) and the American Heart Association (AHA). The most recent version is the 2013 ACCF/AHA Guideline for the Management of Heart Failure [18]. These guidelines were created by a committee of physicians based on thorough review of clinical evidence on heart failure management. They represent a consensus among the physicians on the appropriate treatment and management of heart failure [11]. Though evidence-based guidelines should be the basis for all disease management [6], physicians' adherence to guidelines is very poor [4]. The major reasons for the failure of guideline implementation are lack of awareness, lack of familiarity, lack of motivation and external barriers. For 78% of clinical practice guidelines, more than 10% of the physicians are not aware of their existence. Even when the guidelines are readily accessible, the physicians are not familiar enough with the guidelines to apply them correctly. In all the physician surveys conducted, the lack of familiarity was more common than the lack of awareness [4].

One of the reasons for the lack of familiarity is that the guidelines can be quite complex, as in the case of CHF management. For example, more than 100 variables have been associated with mortality and re-hospitalization related to heart failure. In the 2013 ACCF/AHA Guideline for the Management of Heart Failure, the variables range from simple information like age and sex to sophisticated data like the patterns in electrocardiogram and history of CHF-related symptoms and diseases. To overcome the difficulties that physicians face in implementing the guidelines, we have developed a Physician Advisory System that automates the 2013 ACCF/AHA Guideline for the Management of Heart Failure. Our physician advisory system is able to give recommendations like a real human physician who is following the guidelines strictly, even under the condition of incomplete information about the patient. Our physician-advisory system for CHF management relies on answer set programming [9, 3] for coding the guidelines. The guideline rules are fairly complex and require the use of negation as failure, non-monotonic reasoning and reasoning with incomplete information. A fairly common situation in medicine is that a drug can only be recommended if its use is not contraindicated (i.e., the use of the drug will not have an adverse impact on that patient). Contraindication is naturally modeled via negation as failure. The ability of answer set programming to model defaults, exceptions, weak exceptions, preferences, etc., makes it ideally suited for coding these guidelines.

## **2** Background and overview of the existing literature

A large number of software systems have been designed to address CHF. However, none of them are designed to automatically advise physicians based on the ACCF/AHA guidelines. Chronic disease management systems designed thus far fall into seven categories [12]: accessibility, care management, point-of-care functions, decision support, patient self-management, population management, and reporting. The automation of these functionalities is certainly helpful in assisting health care providers with managing patients with chronic conditions, however, none of them cover what we have realized: a physician advisory system that automates the application of clinical practice guidelines.

The 2013 ACCF/AHA Guideline for the Management of Heart Failure is intended to assist healthcare providers in clinical decision making by describing a range of generally acceptable approaches for the management of chronic heart failure. The guideline is based on four progressive stages of heart failure. Stage A includes patients at risk of heart failure who

are asymptomatic and do not have structural heart disease. Stage B describes asymptomatic patients with structural heart diseases; it includes New York Heart Association (NYHA) class I, in which ordinary physical activity does not cause symptoms of heart failure. Stage C describes patients with structural heart disease who have prior or current symptoms of heart failure; it includes NYHA class I, II (slight limitation of physical activity), III (marked limitation of physical activity) and IV (unable to carry on any physical activity without symptoms of heart failure, or symptoms of heart failure at rest). Stage D describes patients with refractory heart failure who require specialized interventions; it includes NYHA class IV. Interventions at each stage are aimed at reducing risk factors (stage A), treating structural heart disease (stage B) and reducing morbidity and mortality (stages C and D) [18].

Traditional techniques such as logic programming (Prolog) and production systems (OPS5), or traditional expert system styled approaches will result in a far more complex system due to the inability of these systems to model negation as failure effectively [2]. Thus, coding our system in these formalisms would be a much more difficult and complex task. In contrast, the CHF guidelines can be coded in ASP very naturally (it took about 2 months to develop the first version of the system).

### 3 Goal of research

We selected Chronic Heart Failure (CHF) as our first chronic disease to build tools to manage. Chronic Heart Failure is the inability of the heart to pump properly; consequently, not enough oxygen-rich blood can be supplied to all parts of the body. This causes congestion of blood in the lungs, abdomen, legs, etc., causing uneasiness while carrying out any kind of physical activity. Our physician advisory system for CHF management codes all the knowledge in the 2013 ACCF/AHA Guideline for the Management of Heart Failure [18] as an answer set program. Our system is able to recommend treatments just like a human physician who is strictly following these guidelines. Additionally, our system is able to recommend treatments even when a patient's information is incomplete. The input to our system is the patient's information which includes demographics, history, daily symptoms, known risk factors, measurements as well as ACCF/AHA stage and NYHA class [18]. A physician uses our system by posing a query to it. Our system then processes the query by essentially simulating the thinking process of a CHF specialist (represented by the ACCF/AHA guideline).

To implement the CHF guidelines in ASP, we first extensively studied the guidelines to extract reasoning templates. These templates can be thought of as general knowledge patterns. These patterns were next deployed to code the CHF guideline rules. Our research makes two major contributions:

1. We develop a system that completely automates the entire set of guidelines for CHF management developed by the American College of Cardiology Foundation and American Heart Association. The system takes its input from (i) a patient's electronic health record that includes demographic information, test results, etc., and (ii) a telemedicine system that provides data about vital signs (heart rate, blood pressure, weight, etc.). It then uses this information to recommend a treatment. The s(ASP) system also supports abduction, thus our system can also be used for abductive reasoning: a physician can, for example, figure out the symptoms that a particular patient must have in order for a given treatment to work.
2. We develop a set of general knowledge patterns that were used to realize our automated physician-advisory system and that can be helpful in translating rules expressed in a natural language into ASP for any application domain.

## 4 Current status of the research

### 4.1 Physician advisory system description

The physician-advisory system for CHF management has two major components, the rule database and the fact table. The rule database covers all the knowledge in 2013 ACCF/AHA Guideline for the Management of Heart Failure [18]. The fact table contains the relevant information of the patient with heart failure. The fact table is derived from a patient's electronic health record and from a telemedicine system used to measure vital signs. The patient information consists mainly of: 5 pieces of demographics information, 8 measurements and 25 types of HF-related diseases and symptoms. Treatment recommendations returned by the system may include: 11 pharmaceutical treatments, 9 management objectives, and 4 device/surgery therapies.

Our system is designed for running on top of the s(ASP) system, a goal-directed, predicate ASP system that can be thought of as Prolog extended with negation based on the stable model semantics [14]. Because of the goal-directed nature of the system, only the particular treatments applicable to the patient are reported by the system. With minor changes, our system will also work with traditional SAT-based implementations such as CLASP [7, 8]. However, these systems will compute the entire model, so if there are multiple treatments for a given condition, they will all be included in the answer set (these differences between goal-directed and SAT-based solvers are explained in [13]).

### 4.2 Knowledge patterns in the guidelines for the management of heart failure

The ACCF/AHA guidelines are written in English and are quite complex. Our task was to code these guidelines in ASP. To simplify our task, we developed reasoning templates that we call knowledge patterns. These knowledge patterns are quite general and serve as solid building blocks for systematically translating the specifications written in English to ASP. While developing these knowledge patterns and coding them in ASP, certain facts had to be noted: (i) Multiple rules can lead to the recommendation of a treatment; (ii) Multiple rules can lead to contraindication of a treatment; (iii) A treatment cannot be recommended if at least one contraindication for that treatment is present; and, (iv) A given treatment recommendation can impact the recommendation and/or contraindication of other treatments.

Next, we present the most salient knowledge patterns that we have developed. Many of these patterns are straightforward, however, some of them, such as the concomitant choice rule, are intricate. We present these patterns at a high level and ignore non-essential details.

**1. Aggressive Reasoning:** The aggressive reasoning pattern can be stated as “take an action (e.g., recommend treatment) if there is a reason; no evidence of danger means there is no danger in taking that action”. The aggressive reasoning pattern is coded as follows:

```
recommendation(Choice) :- preconditions(Choice),
not contraindication(Choice).
contraindication(Choice) :- dangers(Choice).
```

The code above makes use of negation as failure. If the contraindication of a choice cannot be proved, and the conditions for making the choice hold, then that choice is recommended. An example of this knowledge pattern can be found in [18]: “Digoxin can be beneficial in patients with HF<sub>rEF</sub>, unless contraindicated, to decrease hospitalizations for HF.”

**2. Conservative Reasoning:** This reasoning pattern is stated as “A reason for a recommendation is not enough; evidence that the recommendation is not harmful must be available”.

The conservative reasoning pattern is coded as follows:

```
recommendation(Choice) :- preconditions(Choice),
not contraindication(Choice).
contraindication(Choice) :- not -dangers(Choice).
```

This coding pattern requires evidence of the absence of danger. Without such evidence, the choice would be considered contraindicated. Note that the “-” operator indicates classical negation. An example of this knowledge pattern can be found in [18]: “In patients with structural cardiac abnormalities, including LV hypertrophy, in the absence of a history of MI or ACS, blood pressure should be controlled in accordance with clinical practice guidelines for hypertension to prevent symptomatic HF.”

**3. Anti-recommendation:** The anti-recommendation pattern is stated as “a choice can be prohibited if evidence of danger can be found”.

The coding pattern for the anti-recommendation is coded as follows:

```
contraindication(choice) :- dangers(Choice).
```

The code above specifies the conditions under which a choice will be ruled out (i.e., contraindicated). Note that for a choice to be made, both aggressive reasoning and conservative reasoning require that the contraindication of the choice be false. An example of this knowledge pattern can be found in [18]: “Anticoagulation is not recommended in patients with chronic HFrEF without AF, a prior thromboembolic event, or a cardioembolic source.”

**4. Preference:** The preference pattern is stated as “use the second-line choice when the first-line choice is not available”. The preference pattern is coded as follows:

```
recommendation(First_choice) :- conditions_for_both_choices,
not contraindication(First_choice).
recommendation(Second_choice) :- conditions_for_both_choices,
contraindication(First_choice),
not contraindication(Second_choice).
```

This code chooses the treatment recommendation in the second choice only when the conditions are satisfied, the first choice is contraindicated, and there is no evidence preventing the use of second choice. An example of this knowledge pattern can be found in [18]: “ARBs are recommended in patients with HFrEF with current or prior symptoms who are ACE inhibitor intolerant, unless contraindicated, to reduce morbidity and mortality.”

**5. Concomitant Choice:** The concomitant choice pattern is stated as “if a choice is made, some other choices are automatically in effect unless they are prohibited.” The concomitant pattern is coded as shown below.

```
recommendation(Trigger_choice) :- preconditions(Trigger_choice),
not contraindication(Trigger_choice),
not skip_concomitant_choice(Trigger_choice).
skip_concomitant_choice(Trigger_choice) :-
not recommendation(Concomitant_choice),
```

## 22:6 Automating Disease Management Using Answer Set Programming

```
not contraindication(Concomitant_choice).
recommendation(Concomitant_choice) :-
recommendation(Trigger_choice),
not contraindication(Concomitant_choice).
```

The above code makes sure that a concomitant choice appears in all stable models containing the trigger choice, provided the concomitant choice is not prohibited. The trigger choice is always recommended along with the concomitant choice unless the concomitant choice is contraindicated. An example of this knowledge pattern can be found in [18]: “Diuretics should generally be combined with an ACE inhibitor, beta blocker, and aldosterone antagonist. Few patients with HF will be able to maintain target weight without the use of diuretics.”

**6. Indispensable Choice:** The indispensable choice pattern is stated as “if a choice is made, some other choices must also be made; if those choices can’t be made, then the first choice is revoked”. Note that choosing “Trigger\_choice” forces “Indispensable\_choice”. The indispensable choice pattern is coded as shown below:

```
recommendation(Trigger_choice) :- preconditions(Trigger_choice),
not contraindication(Trigger_choice),
not absent_indispensable_choice(Trigger_choice).
absent_indispensable_choice(Trigger_choice) :-
not recommendation(Indispensable_choice).
recommendation(Indispensable_choice) :- recommendation(Trigger_choice),
not contraindication(Indispensable_choice).
```

The above code makes sure that the trigger choice will always appear with the indispensable choice. If for some reason the indispensable choice cannot be made, then the trigger choice cannot be made either. An example of this knowledge pattern can be found in [18]: “In patients with a current or recent history of fluid retention, beta blockers should not be prescribed without diuretics”.

**7. Incompatible Choice:** The incompatibility pattern is stated as “some choices cannot be in effect at the same time”. The incompatible choice pattern is coded as shown below:

```
taboo_choice(Choice_1) :-
recommendation(Choice_2),
...,
recommendation(Choice_n).
taboo_choice(Choice_2) :-
recommendation(Choice_1),
recommendation(Choice_3),
....
recommendation(Choice_n).
...
taboo_choice(Choice_n) :-
recommendation(Choice_1),
recommendation(Choice_2),
....
recommendation(Choice_n-1).

recommendation(Choice_1) :-
conditions_for_choice_1,
not contraindication(Choice_1),
not taboo_choice(Choice_1).
recommendation(Choice_2) :-
conditions_for_choice_2,
not contraindication(Choice_2),
not taboo_choice(Choice_2).
...
recommendation(Choice_n) :-
conditions_for_choice_n,
not contraindication(Choice_n),
not taboo_choice(Choice_n).
```

```
{accf_stage(c), hf_with_reduced_ef, history(standard_neurohormonal_antagonist_therapy),
nyha_class(3), nyha_class_3_to_4, race(african_american), recommendation(hydralazine_and_isosorbide_dinitrate,class_1),
not contraindication(hydralazine_and_isosorbide_dinitrate)}
```

■ **Figure 1** Result of abductive reasoning in physician-advisory system for CHF management.

The above code makes sure that incompatible choices will not be made together. Note that we did not use a simple constraint to implement this pattern. A constraint would kill all stable models if each of the choices in question can be made. Our implementation, on the other hand, will produce partial answer sets supporting the query, thanks to the goal-driven mechanism of s(ASP) [14]. An example of this knowledge pattern can be found in [18]: “Routine combined use of an ACE inhibitor, ARB, and aldosterone antagonist is potentially harmful for patients with HFrEF.”

### 4.3 Abductive reasoning in the management of heart failure

Our system can also perform abductive reasoning thanks to the s(ASP) system’s support for abduction [14]. Abductive reasoning is a form of logical inference where one attempts to augment a theory with sufficient information to explain an observation (the augmentations come from a set of predicates that are declared as *abducibles*). To illustrate, consider the following two rules in the ACCF/AHA guideline [18]:

- Combination of hydralazine & isosorbide dinitrate is recommended to reduce morbidity & mortality for patients self-described as African Americans with NYHA class III-IV HFrEF receiving optimal therapy with ACE inhibitors & and beta blockers, unless contraindicated.
- Combination of hydralazine & isosorbide dinitrate should not be used for the treatment of HFrEF in patients who have no prior use of standard neurohormonal antagonist therapy.

Suppose we have an African American patient who is suffering from NYHA class III HFrEF, but that is all we know about the patient. Since a hydralazine and isosorbide dinitrate combination is highly effective in reducing the mortality of African Americans with HFrEF, the physician might pose the following query:

```
?-recommendation(hydralazine_and_isosorbide_dinitrate), class_1)
```

to the s(ASP) system. The system would return the results shown in Figure 1.

Note that the system abducted two things: (i) a “history of standard neurohormonal antagonist therapy”, and (ii) the absence of “contraindication of hydralazine and isosorbide dinitrate”. This means in order for us to recommend hydralazine and isosorbide dinitrate to the patient, they must have received standard neurohormonal antagonist therapy before. Otherwise, hydralazine and isosorbide dinitrate would be contraindicated.

## 5 Preliminary results accomplished

Our system has been tested in-house and has shown accurate results that are compatible with what a physician following the guidelines would conclude. A clinical trial is planned.

The input to the system is a patient’s information, including demographics, history, daily symptoms, risks and measurements, as well as ACCF/AHA stage and NYHA class. When queried for a treatment recommendation, our system is able to give recommendations according to the guideline just as a physician would.

```

%doctor's assessments
accf_stage(c).
nyha_class(3).
expectation_of_survival(3).
diagnosis(hypertension).
%demographics of the patient
gender(female).
age(78).
history(mi, recent).
%measurements from the lab
hf_with_reduced_ef.
measurement(creatinine, 1.8).
measurement(potassium, 4.9).
measurement(lvef, 0.35).
measurement(lbbb, 180).
measurement(sinus_rhythm).

%history of the patient
diagnosis(myocardial_ischemia).
diagnosis(atrial_fibrillation).
diagnosis(coronary_artery_disease).

evidence(ischemic_etiology_of_hf).
evidence(sleep_apnea).
evidence(fluid_retention).

history(stroke).
history(cardiovascular_hospitalization).
post_mi(40).

```

■ **Figure 2** Representation of a patient's information in physician-advisory system for CHF management.

To illustrate how our system works, consider a female heart failure patient who is in ACCF/AHA stage C, belongs to NYHA class 3 and has been diagnosed as myocardial ischemia, atrial fibrillation, coronary artery disease. She also suffers from sleep apnea, fluid retention and hypertension. Her left ventricular ejection fraction (LVEF) is 35%. There is evidence that she has ischemic etiology of heart failure. Her electrocardiogram (ECG) has sinus rhythm and a left bundle branch block (LBBB) pattern with a QRS duration of 180ms. The blood test says her creatinine is 1.8 mg/dL and potassium is 4.9 mEq/L. She has a history of stroke. It has been 40 days since the acute myocardial infarction happened to her. Her doctor assessed that her expectation of survival is about 3 years.

The patient's information derived from her electronic health record is coded as the facts shown in Figure 2. There are multiple treatments for this patient. Figure 3 shows some of the treatment recommendations our system infers once we give the query “recommendation(Treatment, Class)”. Each treatment recommendation (represented as a partial answer set) contains all of the predicates that must hold in order for the query to be successful. For instance, consider the recommendation of ace inhibitors as a treatment option (answer #2). Ace inhibitors are recommended because the patient is in ACCF/AHA Stage C, per the doctor's assessment, and has heart failure with reduced ejection fraction condition. Proof of contraindication for ace inhibitors is absent as the patient does not have a history of angioedema (`not history(angioedema)`) and is not pregnant (`not pregnancy`). The system also gives us the concomitant treatments for ace inhibitors, namely, beta blockers and diuretics. It is worth mentioning that we used the aggressive reasoning pattern (see Section 4.2) when coding the rules of ace inhibitors.

Had we adopted the conservative reasoning pattern, ace inhibitors would not have been recommended unless we explicitly asserted `-history(angioedema)` and `-pregnancy` in the patient's information (a definitive proof of the latter can be derived from patient's age (78)).

Given that there may be multiple treatment options for a particular patient, the choice of a particular treatment will depend on the physician's preference. Rules that capture a physician's or a nurse's preference can also be coded as answer set programs in our system.

While our testing indicates that the system works well and the results produced are consistent with what a physician may recommend, if they were to exactly follow the guidelines,

```
{ accf_stage(c), recommendation(sodium_restriction,class_2a), not contraindication(sodium_restriction) } Treatment = sodium_restriction, Class = class_2a
```

```
{ accf_stage(c), hf_with_reduced_ef, recommendation(ace_inhibitors,class_1), recommendation(beta_blockers,class_1), recommendation(diuretics,class_1), not contraindication(ace_inhibitors), not contraindication(beta_blockers), not contraindication(diuretics), not history(angioedema), not history(angioedema,recent), not history(angioedema,remote), not pregnancy } Treatment = ace_inhibitors, Class = class_1
```

■ **Figure 3** Output of the physician-advisory system for CHF management.

a clinical trial is needed to truly validate our system, and is indeed planned. As mentioned earlier, our system can be used for abductive reasoning as well. Running the system in the abductive mode can allow a physician to try out what-if scenarios and to make sure that all the pre-conditions required for treatment are met.

## 6 Open issues and expected achievements

In this paper we report on our work on developing a ASP-based physician advisory system for managing CHF using a telemedicine platform. The system automates the rules laid out in the 2013 ACCF/AHA Guide for the Management of Heart Failure. It is able to take a patient's data as input and produce treatment recommendations that strictly adhere to the guidelines. It can also be used by a physician to abduce symptoms and other conditions that must be met by a given treatment recommendation.

Our approach to developing the system was based on identifying knowledge patterns and using them as building blocks for constructing the ASP code. There are many ways to further extend our work that we plan to pursue in the future:

- Extending the system for comorbidities: We would like our system to handle comorbidities of heart failure [12]. A typical CHF patients suffers from other chronic ailments as well, i.e., CHF generally never occurs by itself.
- Performing clinical trials: our system has been tested in-house, however, we plan to compare the recommendations given by our system to the prescriptions by human cardiologists in a formal clinical trial to validate the effectiveness of our system.
- Integrating with EMRs and a Telemedicine Platform: Future work would include integrating our system with our telemedicine platform so that the input comes directly from the electronic medical record while vital signs are directly obtained from the patient through our telemedicine hardware and software [16, 15]. A user-friendly GUI will also be designed to make the system more usable.
- Adding justification to recommendations given by our system: Although the rationale behind a recommendation is shown in the partial answer set, it is hard to decipher it. We plan to augment s(ASP) [14] so that reasonably detailed justifications for a query are printed in a human-readable form.
- Formal Analysis: Conducting research to formally establish the correctness of our system.

**Acknowledgements.** I want to thank Dr. Gopal Gupta, my supervisor, to introduce me to both logic programming and answer set programming. I would also like to thank Dr. Lakshman Tamil, my co supervisor, for his wonderful insight into chronic disease management. Finally, I want to thank Dr. Kyle Marple and Mr. Elmer Salazar for the enlightening discussions.

---

**References**

---

- 1 Gerard Anderson. *Chronic conditions: making the case for ongoing care*. Johns Hopkins University, 2004.
- 2 Chitta Baral. *Knowledge Representation, Reasoning and Declarative Problem Solving*. Cambridge University Press, 2003.
- 3 Gerhard Brewka, Thomas Eiter, and Mirosław Truszczyński. Answer set programming at a glance. *Commun. ACM* 54, 12, 92–103, 2011.
- 4 Michael D. Cabana, Cynthia S. Rand, Neil R. Powe, and et al. Why don't physicians follow clinical practice guidelines?: A framework for improvement. *JAMA* 282, 15, 1458–1465, 1999.
- 5 Centers for Disease Control and Prevention. Chronic disease overview page. <http://www.cdc.gov/chronicdisease/overview/index.htm>.
- 6 David P. Faxon, et al. Improving quality of care through disease management principles and recommendations from the American heart association's expert panel on disease management. *Stroke* 35, 6, 1527–1530, 2004.
- 7 Martin Gebser, Benjamin Kaufmann, and Torsten Schaub. Conflict-driven answer set solving: From theory to practice. In *Artif. Intell.* 187, 52-89, 2012.
- 8 Martin Gebser, Roland Kaminski, Benjamin Kaufmann, and Torsten Schaub. Clingo = ASP + Control: Preliminary Report. In *Proc. ICLP 2014 Proc.*
- 9 Michael Gelfond and Vladimir Lifschitz. *The stable model semantics for logic programming*. Proc. ICLP 1988. MIT Press, 1070–1080.
- 10 McDonnell Norms Group. Enhancing the use of clinical guidelines: a social norms perspective. *Journal of the American College of Surgeons* 202, 5, 826–836, 2006.
- 11 Alice K. Jacobs, Frederick G. Kushner, and et al. ACCF/AHA clinical practice guideline methodology summit report: A report of the American college of cardiology foundation/american heart association task force on practice guidelines. *Journal of the American College of Cardiology* 61, 2, 213–265, 2013.
- 12 Laura Jantos and Michelle Holmes. *IT Tools for Chronic Disease Management: How Do They Measure Up?* California HealthCare Foundation (chfc.org; retrieved Jan. 2016).
- 13 Kyle Marple and Gopal Gupta. Goal-directed execution of answer set programs. *Proc. PPDP 2012*: 35-44
- 14 Kyle Marple, Elmer Salazar, and Gopal Gupta. s(ASP). <https://sourceforge.net/projects/sasp-system/>.
- 15 Savio Monteiro. An intelligent telemedicine platform with cognitive support for chronic care management. *Ph.D. thesis, Quality of Life Technology Lab, UT Dallas*, 2015.
- 16 Savio Monteiro, Gopal Gupta, Mehrdad Nourani, and Lakshman S Tamil. Hygeiatel: An intelligent telemedicine system with cognitive support. In *Proceedings of the First ACM Workshop on Mobile Systems, Applications, and Services for Healthcare*. mHealthSys'11. ACM, New York, NY, USA, 9:1–2.
- 17 Shin-Yi Wu and Anthony Green. Projection of chronic illness prevalence and cost inflation. *Santa Monica, CA: RAND Health*, 2000.
- 18 Clyde W. Yancy, Mariell Jessup, Biykem Bozkurt, and et al. 2013. 2013 ACCF/AHA guideline for the management of heart failure: A report of the american college of cardiology foundation/American heart association task force on practice guidelines. *Journal of the American College of Cardiology* 62, 16, e147.