

# Separators in Region Intersection Graphs

James R. Lee

Computer Science, University of Washington, Seattle, USA  
jrl@cs.washington.edu

---

## Abstract

For undirected graphs  $G = (V, E)$  and  $G_0 = (V_0, E_0)$ , say that  $G$  is a *region intersection graph over  $G_0$*  if there is a family of connected subsets  $\{R_u \subseteq V_0 : u \in V\}$  of  $G_0$  such that  $\{u, v\} \in E \iff R_u \cap R_v \neq \emptyset$ .

We show if  $G_0$  excludes the complete graph  $K_h$  as a minor for some  $h \geq 1$ , then every region intersection graph  $G$  over  $G_0$  with  $m$  edges has a balanced separator with at most  $c_h \sqrt{m}$  nodes, where  $c_h$  is a constant depending only on  $h$ . If  $G$  additionally has uniformly bounded vertex degrees, then such a separator is found by spectral partitioning.

A string graph is the intersection graph of continuous arcs in the plane. String graphs are precisely region intersection graphs over planar graphs. Thus the preceding result implies that every string graph with  $m$  edges has a balanced separator of size  $O(\sqrt{m})$ . This bound is optimal, as it generalizes the planar separator theorem. It confirms a conjecture of Fox and Pach (2010), and improves over the  $O(\sqrt{m} \log m)$  bound of Matoušek (2013).

**1998 ACM Subject Classification** F.2.2 Nonnumerical Algorithms and Problems, G.1.6 Optimization, G.2.2 Graph Theory

**Keywords and phrases** Graph separators, planar graphs, spectral partitioning

**Digital Object Identifier** 10.4230/LIPIcs.ITCS.2017.1

## 1 Introduction

Consider an undirected graph  $G_0 = (V_0, E_0)$ . A graph  $G = (V, E)$  is said to be a *region intersection graph (rig) over  $G_0$*  if the vertices of  $G$  correspond to connected subsets of  $G_0$  and there is an edge between two vertices of  $G$  precisely when those subsets intersect. Concretely, there is a family of connected subsets  $\{R_u \subseteq V_0 : u \in V\}$  such that  $\{u, v\} \in E \iff R_u \cap R_v \neq \emptyset$ . For succinctness, we will often refer to  $G$  as a *rig over  $G_0$* .

Let  $\text{rig}(G_0)$  denote the family of all finite rigs over  $G_0$ . Prominent examples of such graphs include the intersection graphs of pathwise-connected regions on a surface (which are intersection graphs over graphs that can be drawn on that surface).

For instance, *string graphs* are the intersection graphs of continuous arcs in the plane. It is easy to see that every finite string graph  $G$  is a rig over some planar graph: By a simple compactness argument, we may assume that every two strings intersect a finite number of times. Now consider the planar graph  $G_0$  whose vertices lie at the intersection points of strings and with edges between two vertices that are adjacent on a string (see Figure 1). Then  $G \in \text{rig}(G_0)$ . It is not too difficult to see that the converse is also true; see Section 4.

To illustrate the non-trivial nature of such objects, we recall that there are string graphs on  $n$  strings that require  $2^{\Omega(n)}$  intersections in any such representation [14]. The recognition problem for string graphs is NP-hard [13]. Decidability of the recognition problem was established in [25], and membership in NP was proved in [24]. We refer to the recent survey [22] for more of the background and history behind string graphs.

Even when  $G_0$  is planar, the rigs over  $G_0$  can be dense: Every complete graph is a rig over some planar graph (in particular, every complete graph is a string graph). It has been



© James R. Lee;

licensed under Creative Commons License CC-BY


8th Innovations in Theoretical Computer Science Conference (ITCS 2017).

Editor: Christos H. Papadimitrou; Article No. 1; pp. 1:1–1:8

Leibniz International Proceedings in Informatics



LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany



■ **Figure 1** A string graph as a rig over a planar graph.

conjectured by Fox and Pach [5] that every  $m$ -edge string graph has a balanced separator with  $O(\sqrt{m})$  nodes. Fox and Pach proved that such graphs have separators of size  $O(m^{3/4}\sqrt{\log m})$  and presented a number of applications of their separator theorem. Matoušek [21] obtained a near-optimal bound of  $O(\sqrt{m} \log m)$ . In the present work, we confirm the conjecture of Fox and Pach, and generalize the result to include all rigs over graphs that exclude a fixed minor. This extended abstract contains mostly theorem statements; for detailed proofs and further arguments, we refer to the full paper [15].

► **Theorem 1.** *If  $G \in \text{rig}(G_0)$  and  $G_0$  excludes  $K_h$  as a minor, then  $G$  has a  $\frac{2}{3}$ -balanced separator of size at most  $c_h \sqrt{m}$  where  $m$  is the number of edges in  $G$ . Moreover, one has the estimate  $c_h \leq O(h^3 \sqrt{\log h})$ .*

In the preceding statement, an  $\epsilon$ -balanced separator of  $G = (V, E)$  is a subset  $S \subseteq V$  such that in the induced graph  $G[V \setminus S]$ , every connected component contains at most  $\epsilon|V|$  vertices.

The proof of Theorem 1 is constructive, as it is based on solving and rounding a linear program; it yields a polynomial-time algorithm for constructing the claimed separator. In the case when there is a bound on the maximum degree of  $G$ , one can use the well-known spectral bisection algorithm (see Section 1.5).

Since planar graphs exclude  $K_5$  as a minor, Theorem 1 implies that  $m$ -edge string graphs have  $O(\sqrt{m})$ -node balanced separators. Since the graphs that can be drawn on any compact surface of genus  $g$  exclude a  $K_h$  minor for  $h \leq O(\sqrt{g+1})$ , Theorem 1 also applies to string graphs over any fixed compact surface.

In addition, it implies the Alon-Seymour-Thomas [1] separator theorem<sup>1</sup> for graphs excluding a fixed minor, for the following reason. Let us define the *subdivision* of a graph  $G$  to be the graph  $\dot{G}$  obtained by subdividing every edge of  $G$  into a path of length two. Then every graph  $G$  is a rig over  $\dot{G}$ , and it is not hard to see that for  $h \geq 1$ ,  $G$  has a  $K_h$  minor if and only if  $\dot{G}$  has a  $K_h$  minor.

## 1.1 Applications in topological graph theory

We mention two applications of Theorem 1 in graph theory. In [6], the authors present some applications of separator theorems for string graphs. In two cases, the tight bound for separators leads to tight bounds for other problems. The next two theorems confirm conjectures of Fox and Pach; as proved in [6], they follow from Theorem 1. Both results are tight up to a constant factor.

<sup>1</sup> Note that Theorem 1 is quantitatively weaker in the sense that [1] shows the existence of separators with  $O(h^{3/2}\sqrt{n})$  vertices. Since every  $K_h$ -minor-free graph has at most  $O(nh\sqrt{\log h})$  edges [12, 27], our bound is  $O(h^{7/2}(\log h)^{3/4}\sqrt{n})$ .

► **Theorem 2.** *There is a constant  $c > 0$  such that for every  $t \geq 1$ , it holds that every  $K_{t,t}$ -free string graph on  $n$  vertices has at most  $cnt(\log t)$  edges.*

A *topological graph* is a graph drawn in the plane so that its vertices are represented by points and its edges by curves connecting the corresponding pairs of points.

► **Theorem 3.** *In every topological graph with  $n$  vertices and  $m \geq 4n$  edges, there are two disjoint sets, each of cardinality*

$$\Omega\left(\frac{m^2}{n^2 \log \frac{n}{m}}\right) \tag{1}$$

so that every edge in one set crosses all edges in the other.

This improves over the bound of  $\Omega\left(\frac{m^2}{n^2(\log \frac{n}{m})^c}\right)$  for some  $c > 0$  proved in [7], where the authors also show that the bound (1) is tight. Before we conclude this section, let us justify the observation made earlier.

► **Lemma 4.** *Finite string graphs are precisely finite region intersection graphs over planar graphs.*

**Proof.** We have already argued that string graphs are planar rigs. Consider now a planar graph  $G_0 = (V_0, E_0)$  and a finite graph  $G = (V, E)$  such that  $G \in \text{rig}(G_0)$ . Let  $\{R_u \subseteq V_0 : u \in V\}$  be a representation of  $G$  as a rig over  $G_0$ .

Since  $G$  is finite, we may assume that each region  $R_u$  is finite. To see this, for  $v \in V_0$ , let its *type* be the set  $T(v) = \{u \in V : v \in R_u\}$ . Then since  $G$  is finite, there are only finitely many types. For any region  $R_u \subseteq V_0$ , let  $\tilde{R}_u$  be a finite set of vertices that exhausts every type in  $R_u$ , and let  $\hat{R}_u$  be a finite spanning tree of  $\tilde{R}_u$  in the induced graph  $G_0[R_u]$ . Then the regions  $\{\hat{R}_u : u \in V\}$  are finite and connected, and also form a representation of  $G$  as a rig over  $G_0$ .

When each region  $R_u$  is finite, we may assume also that  $G_0$  is finite. Now take a planar drawing of  $G_0$  in  $\mathbb{R}^2$  where the edges of  $G_0$  are drawn as continuous arcs, and for every  $u \in V$ , let  $T_u \subseteq \mathbb{R}^2$  be the drawing of the spanning tree of  $R_u$ . Each  $T_u$  can be represented by a string (simply trace the tree using an in-order traversal that begins and ends at some fixed node), and thus  $G$  is a string graph. ◀

## 1.2 Balanced separators and extremal spread

Since complete graphs are string graphs, we do not have access to topological methods based on the exclusion of minors. Instead, we highlight a more delicate structural theory. The following fact is an exercise.

**Fact:** If  $\dot{G}$  is a string graph, then  $G$  is planar.

More generally, we recall that  $H$  is a *minor* of  $G$  if  $H$  can be obtained from  $G$  by a sequence of edge contractions, edge deletions, and vertex deletions. If  $H$  can be obtained using only edge contractions and vertex deletions, we say that  $H$  is a *strict minor* of  $G$ .

► **Lemma 5.** *If  $G \in \text{rig}(G_0)$  and  $\dot{H}$  is a strict minor of  $G$ , then  $H$  is a minor of  $G_0$ .*

This topological structure of (forbidden) strict minors in  $G$  interacts nicely with “conformal geometry” on  $G$ . Consider the family of all pseudo-metric spaces that arise from a finite graph  $G$  by assigning non-negative lengths to its edges and taking the induced shortest path

distance. Certainly if we add an edge to  $G$ , the family of such spaces can only grow (since by giving the edge length equal to the diameter of the space, we effectively remove it from consideration). In particular, if  $G = K_n$  is the complete graph on  $n$  vertices, then every  $n$ -point metric space is a path metric on  $G$ .

A significant tool will be the study of extremal conformal metrics on a graph  $G$ . Unlike in the edge-weighted case, the family of path distances coming from conformal metrics can be well-behaved even if  $G$  contains arbitrarily large complete graph minors. As a simple example, let  $K_{\mathbb{N}}$  denote the complete graph on countably many vertices. Then every distance arising from a conformal metric on  $K_{\mathbb{N}}$  is bi-Lipschitz to an ultrametric.

### 1.3 Vertex expansion and observable spread

Fix a graph  $G = (V_G, E_G) \in \text{rig}(G_0)$  with  $n = |V_G|$  and  $m = |E_G|$ . Since the family  $\text{rig}(G_0)$  is closed under taking induced subgraphs, a standard reduction allows us to focus on finding a subset  $U \subseteq V_G$  with small isoperimetric ratio:  $\frac{|\partial U|}{|U|} \lesssim \frac{\sqrt{m}}{n}$ , where

$$\partial U = \{v \in U : E_G(v, V_G \setminus U) \neq \emptyset\},$$

and  $E_G(v, V_G \setminus U)$  is the set of edges between  $v$  and vertices outside  $U$ . Also define the interior  $U^\circ = U \setminus \partial U$ .

Let us define the *vertex expansion constant* of  $G$  as

$$\phi_G = \min \left\{ \frac{|\partial U|}{|U|} : \emptyset \neq U \subseteq V_G, |U^\circ| \leq \frac{|V_G|}{2} \right\}. \quad (2)$$

In [4], it is shown that this quantity is related to the concentration function (in the sense of Lévy and Milman; see also Gromov’s observable diameter [8]) of extremal conformal metrics on  $G$ .

For a finite metric space  $(X, \text{dist})$ , we define the *spread of  $X$*  as the quantity

$$\mathfrak{s}(X, \text{dist}) = \frac{1}{|X|^2} \sum_{x, y \in X} \text{dist}(x, y).$$

Define the *observable spread of  $X$*  by

$$\mathfrak{s}_{\text{obs}}(X, \text{dist}) = \sup_{f: X \rightarrow \mathbb{R}} \left\{ \frac{1}{|X|^2} \sum_{x, y \in X} |f(x) - f(y)| : f \text{ is 1-Lipschitz} \right\}. \quad (3)$$

► **Remark.** We remark on the terminology: In general, it is difficult to “view” a large metric space all at once; this holds both conceptually and from an algorithmic standpoint. If one thinks of Lipschitz maps  $f : X \rightarrow \mathbb{R}$  as “observations” then the observable spread captures how much of the spread can be “seen.”

We then define the  *$L^1$ -extremal observable spread of  $G$*  as

$$\bar{\mathfrak{s}}_{\text{obs}}(G) = \sup_{\omega: V_G \rightarrow \mathbb{R}_+} \left\{ \mathfrak{s}_{\text{obs}}(V_G, \text{dist}_\omega) : \|\omega\|_{L^1(V_G)} \leq 1 \right\}, \quad (4)$$

where  $\|\omega\|_{L^1(V_G)} := \frac{1}{|V_G|} \sum_{v \in V_G} \omega(v)$ . We recall the following theorem from [4] that relates expansion to the observable spread.

► **Theorem 6 ([4]).** *For every finite graph  $G$ ,*

$$\frac{1}{2} \bar{\mathfrak{s}}_{\text{obs}}(G) \leq \frac{1}{\phi_G} \leq 3 \bar{\mathfrak{s}}_{\text{obs}}(G).$$

► **Example 7.** If  $G$  is the subgraph of the lattice  $\mathbb{Z}^d$  on the vertex set  $\{0, 1, \dots, L\}^d$ , then  $\phi_G \asymp 1/L$  and  $\bar{\mathfrak{s}}(G) \asymp L$ . This can be achieved by taking  $\omega \equiv 1$  and defining  $f : V_G \rightarrow \mathbb{R}$  by  $f(x) = x_1$ .

In light of Theorem 6, to prove Theorem 1, it suffices to give a lower bound on  $\bar{\mathfrak{s}}_{\text{obs}}(G)$ . It is natural to compare this quantity to the  $L^1$ -extremal spread of  $G$ :

$$\bar{\mathfrak{s}}(G) := \max \left\{ \frac{1}{|V_G|^2} \sum_{u,v \in V_G} \text{dist}_\omega(u,v) : \|\omega\|_{L^1(V_G)} \leq 1 \right\}. \quad (5)$$

Let us examine these two notions for planar graphs using the theory of circle packings.

► **Example 8 (Circle packings).** Suppose that  $G$  is a finite planar graph. The Koebe-Andreiev-Thurston circle packing theorem asserts that  $G$  is the tangency graph of a family  $\{D_v : v \in V_G\}$  of circles on the unit sphere  $\mathbb{S}^2 \subseteq \mathbb{R}^3$ . Let  $\{c_v : v \in V_G\} \subseteq \mathbb{S}^2$  and  $\{r_v > 0 : v \in V_G\}$  be the centers and radii of the circles, respectively. An argument of Spielman and Teng [26] (see also Hersch [9] for the analogous result for conformal mappings) shows that one can take  $\sum_{v \in V_G} c_v = \mathbf{0}$ .

If we define  $\omega(v) = r_v$  for  $v \in V_G$ , then  $\text{dist}_\omega \geq \text{dist}_{\mathbb{S}^2} \geq \text{dist}_{\mathbb{R}^3}$  on the centers  $\{c_v : v \in V_G\}$ . (The latter two distances are the geodesic distance on  $\mathbb{S}^2$  and the Euclidean distance on  $\mathbb{R}^3$ , respectively).

Using the fact that  $\sum_{v \in V_G} c_v = \mathbf{0}$ , we have

$$\sum_{u,v \in V_G} \|c_u - c_v\|_2^2 = 2n \sum_{u \in V_G} \|c_u\|^2 = 2n^2. \quad (6)$$

This yields

$$\sum_{u,v \in V_G} \text{dist}_\omega(u,v) \geq \sum_{u,v \in V_G} \|c_u - c_v\| \geq \frac{n^2}{2}.$$

Moreover,

$$\|\omega\|_{L^1(V_G)} \leq \|\omega\|_{L^2(V_G)} = \sqrt{\frac{1}{n} \sum_{v \in V_G} r_v^2} \leq \sqrt{\frac{\text{vol}(\mathbb{S}^2)}{\pi n}} = \sqrt{\frac{4}{n}}.$$

It follows that  $\bar{\mathfrak{s}}(G) \geq \frac{\sqrt{n}}{4}$ .

Observe that the three coordinate projections  $\mathbb{R}^3 \rightarrow \mathbb{R}$  are all Lipschitz with respect to  $\text{dist}_\omega$ , and one of them contributes at least a 1/3 fraction to the sum (6). We conclude that  $\bar{\mathfrak{s}}_{\text{obs}}(G) \geq \frac{\sqrt{n}}{12}$ . Combined with Theorem 6, this yields a proof of the Lipton-Tarjan separator theorem [18]. Similar proofs of the separator theorem based on circle packings are known (see [23]), and this one is not new (certainly it was known to the authors of [26]).

We will prove Theorem 1 in two steps: By first giving a lower bound  $\bar{\mathfrak{s}}(G) \gtrsim n/\sqrt{m}$  and then establishing  $\bar{\mathfrak{s}}_{\text{obs}}(G) \gtrsim \bar{\mathfrak{s}}(G)$ .

For the first step, we follow [21, 4, 2]. The optimization (5) is a linear program, and the dual optimization is a maximum multi-flow problem in  $G$ . Matoušek shows that a low-congestion multi-flow can be used to draw the complete graph in the plane with few edge crossings. Since this is impossible by a simple double-counting argument, one concludes that there is no low-congestion flow, providing a lower bound on  $\bar{\mathfrak{s}}(G)$  via LP duality. We extend this argument to rigs over  $K_h$ -minor-free graphs using the flow crossing framework of [2].

### 1.4 Spread vs. observable spread

Our major departure from [21] comes in the second step: Rounding a fractional separator to an integral separator by establishing that  $\bar{s}_{\text{obs}}(G) \geq C_h \cdot \bar{s}(G)$  when  $G$  is a rig over a  $K_h$ -minor-free graph. Matoušek used the following result that holds for any metric space. It follows easily from the methods of [3] or [17] (see also [20, Ch. 15]).

► **Theorem 9.** *For any finite metric space  $(X, d)$  with  $|X| \geq 2$ , it holds that*

$$s_{\text{obs}}(X, d) \geq \frac{s(X, d)}{O(\log |X|)}.$$

In particular, for any graph  $G$  on  $n \geq 2$  vertices,

$$\bar{s}_{\text{obs}}(G) \geq \frac{\bar{s}(G)}{O(\log n)}.$$

Instead of using the preceding result, we employ the graph partitioning method of Klein, Plotkin, and Rao [11]. Those authors present an iterative process for repeatedly partitioning a metric graph  $G$  until the diameter of the remaining components is bounded. If the partitioning process fails, they construct a  $K_h$  minor in  $G$ .

Since rigs over  $K_h$ -minor-free graphs do not necessarily exclude any minors, we need to construct a different sort of forbidden structure. This is the role that Lemma 5 plays in [15]. In order for the argument to work, it is essential that we construct induced partitions: We remove a subset of the vertices which induces a partitioning of the remainder into connected components. After constructing a suitable random partition of  $G$ , standard methods from metric embedding theory allow us to conclude

### 1.5 Eigenvalues and $L^2$ -extremal spread

The methods presented here can be used to control eigenvalues of the discrete Laplacian on rigs. Consider the linear space  $\mathbb{R}^{V_G} = \{f : V_G \rightarrow \mathbb{R}\}$ . Let  $\mathcal{L}_G : \mathbb{R}^{V_G} \rightarrow \mathbb{R}^{V_G}$  be the symmetric, positive semi-definite linear operator given by

$$\mathcal{L}_G f(v) = \sum_{u:\{u,v\} \in E_G} (f(v) - f(u)).$$

Let  $0 = \lambda_0(G) \leq \lambda_1(G) \leq \dots \leq \lambda_{|V_G|-1}(G)$  denote the spectrum of  $\mathcal{L}_G$ .

Define the  $L^p$ -extremal spread of  $G$  as

$$\bar{s}_p(G) = \max_{\omega: V_G \rightarrow \mathbb{R}_+} \left\{ \frac{1}{|V_G|^2} \sum_{u,v \in V_G} \text{dist}_\omega(u, v) : \|\omega\|_{L^p(V_G)} \leq 1 \right\}. \tag{7}$$

In [2], the  $L^2$ -extremal spread is used to give upper bounds on the first non-trivial eigenvalue of graphs that exclude a fixed minor. In [10], a stronger property of conformal metrics is used to bound the higher eigenvalues as well. Roughly speaking, to control the  $k$ th eigenvalue, one requires a conformal metric  $\omega : V_G \rightarrow \mathbb{R}_+$  such that the spread on every subset of size  $\geq |V_G|/k$  remains large. Combining their main theorems with our methods yields the following.

► **Theorem 10.** *Suppose that  $G \in \text{rig}(G_0)$  and  $G_0$  excludes  $K_h$  as a minor for some  $h \geq 3$ . If  $d_{\text{max}}$  is the maximum degree of  $G$ , then for any  $k = 1, 2, \dots, |V_G| - 1$ , it holds that*

$$\lambda_k(G) \leq O(d_{\text{max}}^2 h^6 \log h) \frac{k}{|V_G|}.$$

In particular, the bound on  $\lambda_1(G)$  shows that if  $d_{\max}(G) \leq O(1)$ , then recursive spectral partitioning (see [26]) finds an  $O(\sqrt{n})$ -vertex balanced separator in  $G$ .

## 1.6 Additional applications

**Treewidth approximations.** Bounding  $\bar{s}_{\text{obs}}(G)$  for rigs over  $K_h$ -minor-free graphs leads to some additional applications. Combined with the rounding algorithm implicit in Theorem 6 (and explicit in [4]), this yields an  $O(h^2)$ -approximation algorithms for the vertex uniform Sparsest Cut problem. In particular, it follows that if  $G \in \text{rig}(G_0)$  and  $G_0$  excludes  $K_h$  as a minor, then there is a polynomial-time algorithm that constructs a tree decomposition of  $G$  with treewidth  $O(h^2 \text{tw}(G))$ , where  $\text{tw}(G)$  is the treewidth of  $G$ . This result appears new even for string graphs. We refer to [4].

**Lipschitz extension.** Our results on padded decomposability of conformal metrics on string graphs combine with the Lipschitz extension theory of [16] to show the following. Suppose that  $(G, \omega)$  is a conformal graph, where  $G$  is a rig over some  $K_h$ -minor free graph. Then for every Banach space  $Z$ , subset  $S \subseteq V_G$ , and  $L$ -Lipschitz mapping  $f : S \rightarrow Z$ , there is an  $O(h^2 L)$ -Lipschitz extension  $\tilde{f} : V_G \rightarrow Z$  with  $\tilde{f}|_S = f$ . See [19] for applications to flow and cut sparsifiers in such graphs.

**Acknowledgements.** The author thanks Noga Alon, Nati Linial, and Laci Lovász for helpful discussions, Janos Pach for emphasizing Jirka's near-optimal bound for separators in string graphs, and the organizers of the "Mathematics of Jiří Matoušek" conference, where much of this work was carried out.

---

## References

- 1 Noga Alon, Paul Seymour, and Robin Thomas. A separator theorem for nonplanar graphs. *J. Amer. Math. Soc.*, 3(4):801–808, 1990.
- 2 Punyashloka Biswal, James R. Lee, and Satish Rao. Eigenvalue bounds, spectral partitioning, and metrical deformations via flows. *J. ACM*, 57(3):Art. 13, 23, 2010. Prelim. version in *FOCS 2008*. doi:10.1145/1706591.1706593.
- 3 J. Bourgain. On Lipschitz embedding of finite metric spaces in Hilbert space. *Israel J. Math.*, 52(1-2):46–52, 1985.
- 4 Uriel Feige, MohammadTaghi Hajiaghayi, and James R. Lee. Improved approximation algorithms for minimum weight vertex separators. *SIAM J. Comput.*, 38(2):629–657, 2008. doi:10.1137/05064299X.
- 5 Jacob Fox and János Pach. A separator theorem for string graphs and its applications. *Combin. Probab. Comput.*, 19(3):371–390, 2010. doi:10.1017/S0963548309990459.
- 6 Jacob Fox and János Pach. Applications of a new separator theorem for string graphs. *Combin. Probab. Comput.*, 23(1):66–74, 2014. doi:10.1017/S0963548313000412.
- 7 Jacob Fox, János Pach, and Csaba D. Tóth. A bipartite strengthening of the crossing lemma. *J. Combin. Theory Ser. B*, 100(1):23–35, 2010. doi:10.1016/j.jctb.2009.03.005.
- 8 Misha Gromov. *Metric structures for Riemannian and non-Riemannian spaces*. Modern Birkhäuser Classics. Birkhäuser Boston, Inc., Boston, MA, english edition, 2007. Based on the 1981 French original, With appendices by M. Katz, P. Pansu and S. Semmes, Translated from the French by Sean Michael Bates.
- 9 Joseph Hersch. Quatre propriétés isopérimétriques de membranes sphériques homogènes. *C. R. Acad. Sci. Paris Sér. A-B*, 270:A1645–A1648, 1970.

- 10 J. Kelner, J. R. Lee, G. Price, and S.-H. Teng. Metric uniformization and spectral bounds for graphs. *Geom. Funct. Anal.*, 21(5):1117–1143, 2011. Prelim. version in *STOC 2009*.
- 11 Philip N. Klein, Serge A. Plotkin, and Satish Rao. Excluded minors, network decomposition, and multicommodity flow. In *Proceedings of the 25th Annual ACM Symposium on Theory of Computing*, pages 682–690, 1993.
- 12 A. V. Kostochka. The minimum Hadwiger number for graphs with a given mean degree of vertices. *Metody Diskret. Analiz.*, (38):37–58, 1982.
- 13 Jan Kratochvíl. String graphs. II. Recognizing string graphs is NP-hard. *J. Combin. Theory Ser. B*, 52(1):67–78, 1991. doi:10.1016/0095-8956(91)90091-W.
- 14 Jan Kratochvíl and Jiří Matoušek. String graphs requiring exponential representations. *J. Combin. Theory Ser. B*, 53(1):1–4, 1991. doi:10.1016/0095-8956(91)90050-T.
- 15 James R. Lee. Separators in region intersection graphs. Available at arXiv:math/1608.01612, 2016.
- 16 James R. Lee and Assaf Naor. Extending Lipschitz functions via random metric partitions. *Invent. Math.*, 160(1):59–95, 2005.
- 17 Tom Leighton and Satish Rao. Multicommodity max-flow min-cut theorems and their use in designing approximation algorithms. *J. ACM*, 46(6):787–832, 1999.
- 18 Richard J. Lipton and Robert Endre Tarjan. A separator theorem for planar graphs. *SIAM J. Appl. Math.*, 36(2):177–189, 1979. doi:10.1137/0136016.
- 19 Konstantin Makarychev and Yury Makarychev. Metric extension operators, vertex sparsifiers and Lipschitz extendability. *Israel J. Math.*, 212(2):913–959, 2016. doi:10.1007/s11856-016-1315-8.
- 20 J. Matoušek. *Lectures on discrete geometry*, volume 212 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2002.
- 21 Jiří Matoušek. Near-optimal separators in string graphs. *Combin. Probab. Comput.*, 23(1):135–139, 2014. doi:10.1017/S0963548313000400.
- 22 Jiří Matoušek. String graphs and separators. In *Geometry, structure and randomness in combinatorics*, volume 18 of *CRM Series*, pages 61–97. Ed. Norm., Pisa, 2015.
- 23 Gary L. Miller, Shang-Hua Teng, William Thurston, and Stephen A. Vavasis. Separators for sphere-packings and nearest neighbor graphs. *J. ACM*, 44(1):1–29, 1997. doi:10.1145/256292.256294.
- 24 Marcus Schaefer, Eric Sedgwick, and Daniel Štefankovič. Recognizing string graphs in NP. *J. Comput. System Sci.*, 67(2):365–380, 2003. Special issue on STOC2002 (Montreal, QC). doi:10.1016/S0022-0000(03)00045-X.
- 25 Marcus Schaefer and Daniel Štefankovič. Decidability of string graphs. *J. Comput. System Sci.*, 68(2):319–334, 2004. doi:10.1016/j.jcss.2003.07.002.
- 26 Daniel A. Spielman and Shang-Hua Teng. Spectral partitioning works: Planar graphs and finite element meshes. *Linear Algebra and its Applications: Special Issue in honor of Miroslav Fiedler*, 421(2–3):284–305, March 2007.
- 27 Andrew Thomason. An extremal function for contractions of graphs. *Math. Proc. Cambridge Philos. Soc.*, 95(2):261–265, 1984.