

Tensor of Quantitative Equational Theories

Giorgio Bacci  

Department of Computer Science, Aalborg University, Denmark

Radu Mardare 

Department of Computer & Information Sciences, University of Strathclyde, Glasgow, UK

Prakash Panangaden 

School of Computer Science, McGill University, Montreal, Canada

Gordon Plotkin 

LFCS, School of Informatics, University of Edinburgh, UK

Abstract

We develop a theory for the commutative combination of quantitative effects, their *tensor*, given as a combination of quantitative equational theories that imposes mutual commutation of the operations from each theory. As such, it extends the sum of two theories, which is just their unrestrained combination. Tensors of theories arise in several contexts; in particular, in the semantics of programming languages, the monad transformer for global state is given by a tensor.

We show that under certain assumptions on the quantitative theories the free monad that arises from the tensor of two theories is the categorical tensor of the free monads on the theories. As an application, we provide the first algebraic axiomatizations of labelled Markov processes and Markov decision processes. Apart from the intrinsic interest in the axiomatizations, it is pleasing they are obtained compositionally by means of the sum and tensor of simpler quantitative equational theories.

2012 ACM Subject Classification Theory of computation → Equational logic and rewriting

Keywords and phrases Quantitative equational theories, Tensor, Monads, Quantitative Effects

Digital Object Identifier 10.4230/LIPIcs.CALCO.2021.7

1 Introduction

The theory of computational effects began with the work of Moggi [24, 25] seeking a unified category-theoretic account of the semantics of higher-order programming languages. He modelled computational effects (which he called notions of computation) by means of strong monads on a base category with cartesian closed structure. With Cenciarelli [5], he later extended the theory by allowing a compositional treatment of various semantic phenomena such as state, IO, exceptions, resumptions, etc, via the use of monad transformers. This work was followed up by the program of Plotkin and Power [26, 27] on an axiomatic understanding of computational effects as arising from operations and equations via the use of Lawvere theories (see also [14]). In a fundamental contribution [12] together with Hyland they developed a unified modular theory for algebraic effects that supports their combination by taking the *sum* and *tensor* of their Lawvere theories. This allowed them to recover in a more pleasing algebraic structural way many of the monad transformers considered by Moggi.

Quantitative equational theories, introduced by Mardare et al. [21], are a logical framework generalising the standard concept of equational logic to account for a concept of approximate equality. They are used to describe quantitative effects as monads on categories of metric spaces. Following the work of Hyland et al. [12], in [1] we developed a theory for the sum of quantitative equational theories, and proved that it corresponds to the categorical sum of quantitative effects as monads. As a major example, we axiomatised Markov processes with discounted probabilistic bisimilarity distance [7] as the sum of two theories: interpolative barycentric algebras (which axiomatises probability distributions with the Kantorovich metric [21]) and contractive operators (used to express the transition to the next state).



© Giorgio Bacci, Radu Mardare, Prakash Panangaden, and Gordon Plotkin;
licensed under Creative Commons License CC-BY 4.0

9th Conference on Algebra and Coalgebra in Computer Science (CALCO 2021).

Editors: Fabio Gadducci and Alexandra Silva; Article No. 7; pp. 7:1–7:17

Leibniz International Proceedings in Informatics



LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

Whereas the sum of two monads is the simplest combination supporting both given effects with no interactions between them, the tensor additionally requires commutation of these effects over each other. Some of the most important monad transformers have an elegant abstract description using tensor. Specifically, Moggi’s transformers for state, reader, and writer are examples of tensors [12].

In the present paper we extend the work initiated in [1], and develop the theory for the *tensor of quantitative equational theories*. The main contributions are:

1. we prove that the tensor of quantitative theories corresponds to the categorical tensor of their induced quantitative effects as strong monads;
2. we give quantitative axiomatisations to the *quantitative reader* and *writer monads*, from which we obtain analogues of Moggi’s transformers at the level of theories using tensor;
3. we provide the first axiomatization of *labelled Markov processes* and *Markov decision processes* with their discounted bisimilarity metrics.

For the proof of (1) we introduce the concept of *pre-operation of a strong functor*, which we use to conveniently characterise the commutative bialgebras for the monads (which correspond to the Eilenberg-Moore algebras for their tensor). Crucially, this allows us to carry out the technical development directly at the level of quantitative equational theories without passing via a correspondence with metric-enriched Lawvere theories.

The axiomatisations in (3) are two major examples for our compositional theory quantitative effects. Specifically, we obtain the discounted bisimilarity metrics for labelled Markov processes and Markov decision processes with rewards by complementing the axiomatization for Markov processes presented in [1]. We model reactions to action labels by tensoring with the theory of quantitative reading computations (corresponding to Moggi’s reader monad transformer); while rewards are recovered by tensoring with the theory of quantitative writing computations (corresponding to Moggi’s writer monad transformer). We will illustrate our compositional approach by decomposing the proposed axiomatisations into their basic components and showing how to combine them step-by-step to get the desired result.

Further Related Work. In [12, 11] the tensor of (enriched) Lawvere theories is characterized as the colimit of certain commutative cocones, and the correspondence with the tensor of monads is obtained via the equivalence between Lawvere theories and monads. Since it is not hard to show that (basic) quantitative equational theories can be characterised as metric-enriched Lawvere theories, one may think to recover the correspondence with the tensor of monads via the equivalence with Lawvere theories. Alas, quantitative equational theories and Lawvere theories are not equivalent, as the latter allows generic operations with metric spaces as arities, while the framework of Mardare et al. [21] does not. An equivalence with *discrete* Lawvere theories [13] (where arities are just countable ordinals) does not hold either, because quantitative equations implicitly impose the existence operations with non-discrete arities which cannot be expressed in the framework of discrete Lawvere theories.

The above arguments required us to follow a different path, which led us to the introduction of pre-operations for a strong functor F . Pre-operations are related to Plokin and Power’s *algebraic operations* [28, 29] in the sense that their assignment to F -algebras are the appropriate version of algebraic operations for functors. Moreover, when considered over a strong monad T they correspond to generic effects of type $I \rightarrow Tv$ (*i.e.*, Kleisli maps of type $I \rightarrow v$, where I is the identity for the monoidal product). The reason why we consider pre-operations over functors, and not just monads, is to relate the operations of an algebraic monad with those of its signature. This was crucial in the technical development of Section 5.

Finally, we remark that quantitative equational theories are a natural kind of enriched equational theory expressive enough to recover many examples of interest in computer science (see [21, 1, 23]), but not corresponding to metric-enriched Lawvere theories. In this respect, it is nice that also this simpler subclass of enriched theories are closed under sum and tensor.

2 Preliminaries and Notation

An *extended metric space* is a pair (X, d) consisting of a set X equipped with a distance function $d: X \times X \rightarrow [0, \infty]$ allowed to have infinite values, satisfying: (i) $d(x, y) = 0$ iff $x = y$, (ii) $d(x, y) = d(y, x)$ and (iii) $d(x, z) \leq d(x, y) + d(y, z)$.

A sequence (x_i) in X is *Cauchy* if $\forall \epsilon > 0, \exists N, \forall i, j \geq N, d(x_i, x_j) \leq \epsilon$. If every Cauchy sequence converges, the extended metric space (X, d) is said to be *complete*. If a space is not complete it can be completed by a well-known construction called *Cauchy completion*. We write $\overline{(X, d)}$, or just \overline{X} , for the completion of (X, d) .

We denote by **Met** the category of extended metric spaces with morphisms the non-expansive maps, *i.e.* the $f: (X, d_X) \rightarrow (Y, d_Y)$ such that $d_X(x, y) \geq d_Y(f(x), f(y))$. This category is both complete (*i.e.*, have all limits) and cocomplete (*i.e.*, have all colimits). We will consider also the full subcategory **CMet** of complete extended metric spaces.

The categorical properties of extended metric spaces are much nicer than usual metric spaces. In particular, we note that **Met** is a symmetric monoidal category, with monoidal product $(X, d_X) \square (Y, d_Y)$ being the extended metric space with underlying set $X \times Y$ and extended metric $d_{X \square Y}((x, y), (x', y')) = d_X(x, x') + d_Y(y, y')$ (*cf.* [19]). Note that this is not the cartesian product in **Met** (for which $+$ above would be replaced by \max).

The monoidal product \square introduced above defines a closed monoidal structure on **Met**, with internal hom $[(X, d_X), (Y, d_Y)]$ given by the set of non-expansive maps from X to Y with point-wise supremum extended metric $d_{[(X, d_X), (Y, d_Y)]}(f, g) = \sup_{x \in X} d(f(x), g(x))$.

Finally, we recall the basic definitions of strong functor (and monad), strong natural transformations, and fix the notation (for more details see e.g. [17, 18]). Let \mathbf{V} be a symmetric monoidal closed category with monoidal product¹ $\square: \mathbf{V} \times \mathbf{V} \rightarrow \mathbf{V}$, unit object $I \in \mathbf{V}$, and internal hom-functor $[-, -]: \mathbf{V} \times \mathbf{V} \rightarrow \mathbf{V}$. We will denote the *counit* (or evaluation map) of the adjunction $(V \square -) \dashv [V, -]$ by $ev^V: V \square [V, -] \Rightarrow Id$ and the *unit* (or co-evaluation map) by $\overline{ev}^V: Id \rightarrow [V, V \square -]$.

A functor $F: \mathbf{V} \rightarrow \mathbf{V}$ is *strong* with *monoidal strength* $t_{V, W}: V \square F(W) \rightarrow F(V \square W)$, if t is a natural transformation satisfying the coherence conditions $F\lambda \circ t = \lambda$ and $t \circ (id \square t) \circ \alpha = F\alpha \circ t$, w.r.t. the associator α and left unitor λ of \mathbf{V} . The dual strength $\hat{t}_{V, W}: F(W) \square V \rightarrow F(W \square V)$ is given by $\hat{t} = F(s) \circ t \circ s$, where $s: V \square W \rightarrow W \square V$ is the natural isomorphism of the symmetric monoidal category \mathbf{V} . A natural transformation $\theta: F \Rightarrow G$ is said *strong* if F, G are strong functors with strengths t, σ , respectively, and $\sigma \circ (id \square \theta) = \theta \circ t$, meaning that θ interacts well with the strengths.

A monad (T, η, μ) with unit $\eta: Id \Rightarrow T$ and multiplication $\mu: TT \Rightarrow T$, is *strong* if T is a strong functor with strength t such that $t \circ (id \square \eta) = \eta$ and $\mu \circ tt = t \circ (id \square \mu)$.

Note that strong functors (resp. monads) on a symmetric monoidal closed category \mathbf{V} are equivalent to \mathbf{V} -enriched functors (resp. monads) on the self-enriched category \mathbf{V} [17].

3 Quantitative Equational Theories

Quantitative equations were introduced in [21]. In this framework equalities $t \equiv_\epsilon s$ are indexed by a positive rational number, to capture the idea that t is “within ϵ ” of s . This informal notion is formalised in a manner analogous to traditional equational logic. In this section we review this formalism.

¹ We denote the monoidal product by \square to avoid confusion with other tensorial operations we will deal with in this paper, e.g., the tensor of monads.

7:4 Tensor of Quantitative Equational Theories

Let Σ be a signature of function symbols $f: n \in \Sigma$ of arity $n \in \mathbb{N}$. Let X be a countable set of variables, ranged over by x, y, z, \dots . We write $\mathbb{T}(\Sigma, X)$ for the set of Σ -terms freely generated over X , ranged over by t, s, u, \dots .

A *substitution of type Σ* is a function $\sigma: X \rightarrow \mathbb{T}(\Sigma, X)$, canonically extended to terms as $\sigma(f(t_1, \dots, t_n)) = f(\sigma(t_1), \dots, \sigma(t_n))$; we write $\mathcal{S}(\Sigma)$ for the set of substitutions of type Σ .

A *quantitative equation of type Σ over X* is an expression of the form $t \equiv_{\varepsilon} s$, for $t, s \in \mathbb{T}(\Sigma, X)$ and $\varepsilon \in \mathbb{Q}_{\geq 0}$. We use $\mathcal{V}(\Sigma, X)$ to denote the set of quantitative equations of type Σ over X , and its subsets will be ranged over by Γ, Θ, \dots . Let $\mathcal{E}(\Sigma, X)$ be the set of *conditional quantitative equations* on $\mathbb{T}(\Sigma, X)$, which are expressions of the form

$$\{t_1 \equiv_{\varepsilon_1} s_1, \dots, t_n \equiv_{\varepsilon_n} s_n\} \vdash t \equiv_{\varepsilon} s,$$

for arbitrary $s_i, t_i, s, t \in \mathbb{T}(\Sigma, X)$ and $\varepsilon_i, \varepsilon \in \mathbb{Q}_{\geq 0}$.

A *quantitative equational theory of type Σ over X* is a set \mathcal{U} of conditional quantitative equations on $\mathbb{T}(\Sigma, X)$ containing the axioms and closed under the rules of inference below, for arbitrary $x, y, z, x_i, y_i \in X$, terms $s, t \in \mathbb{T}(\Sigma, X)$, rationals $\varepsilon, \varepsilon' \in \mathbb{Q}_{\geq 0}$, and $\Gamma, \Theta \subseteq \mathcal{V}(\Sigma, X)$,

- (Refl) $\vdash x \equiv_0 x$,
- (Symm) $\{x \equiv_{\varepsilon} y\} \vdash y \equiv_{\varepsilon} x$,
- (Triang) $\{x \equiv_{\varepsilon} z, z \equiv_{\varepsilon'} y\} \vdash x \equiv_{\varepsilon+\varepsilon'} y$,
- (Max) $\{x \equiv_{\varepsilon} y\} \vdash x \equiv_{\varepsilon+\varepsilon'} y$, for all $\varepsilon' > 0$,
- (Cont) $\{x \equiv_{\varepsilon'} y \mid \varepsilon' > \varepsilon\} \vdash x \equiv_{\varepsilon} y$,
- (f -NE) $\{x_i \equiv_{\varepsilon} y_i \mid i=1..n\} \vdash f(x_1, \dots, x_n) \equiv_{\varepsilon} f(y_1, \dots, y_n)$, for $f: n \in \Sigma$,
- (Subst) If $\Gamma \vdash t \equiv_{\varepsilon} s$, then $\{\sigma(t) \equiv_{\varepsilon} \sigma(s) \mid t \equiv_{\varepsilon} s \in \Gamma\} \vdash \sigma(t) \equiv_{\varepsilon} \sigma(s)$, for $\sigma \in \mathcal{S}(\Sigma)$,
- (Ass) If $t \equiv_{\varepsilon} s \in \Gamma$, then $\Gamma \vdash t \equiv_{\varepsilon} s$,
- (Cut) If $\Gamma \vdash \Theta$ and $\Theta \vdash t \equiv_{\varepsilon} s$, then $\Gamma \vdash t \equiv_{\varepsilon} s$,

where we write $\Gamma \vdash \Theta$ to mean that $\Gamma \vdash t \equiv_{\varepsilon} s$ holds for all $t \equiv_{\varepsilon} s \in \Theta$.

The rules (Subst), (Cut), (Ass) are the usual rules of equational logic. The axioms (Refl), (Symm), (Triang) correspond, respectively, to reflexivity, symmetry, and the triangle inequality; (Max) represents inclusion of neighborhoods of increasing diameter; (Cont) is the limiting property of a decreasing chain of neighborhoods with converging diameters; and (f -NE) expresses non-expansiveness of $f \in \Sigma$.

A set A of conditional quantitative equations *axiomatizes* a quantitative equational theory \mathcal{U} , if \mathcal{U} is the smallest quantitative equational theory containing A .

The models of these theories, called *quantitative Σ -algebras*, are Σ -algebras in **Met**.

► **Definition 1** (Quantitative Algebra). A quantitative Σ -algebra is a tuple $\mathcal{A} = (A, \Sigma^{\mathcal{A}})$, where A is an extended metric space and $\Sigma^{\mathcal{A}} = \{f^{\mathcal{A}}: A^n \rightarrow A \mid f: n \in \Sigma\}$ is a set of non-expansive interpretations (i.e., satisfying $\max_i d_A(a_i, b_i) \geq d_A(f^{\mathcal{A}}(a_1, \dots, a_n), f^{\mathcal{A}}(b_1, \dots, b_n))$).

The morphisms between quantitative Σ -algebras are non-expansive Σ -homomorphisms. Quantitative Σ -algebras and their morphism form a category, denoted by **QA**(Σ).

$\mathcal{A} = (A, \Sigma^{\mathcal{A}})$ satisfies the conditional quantitative equation $\Gamma \vdash t \equiv_{\varepsilon} s$ in $\mathcal{E}(\Sigma, X)$, written $\Gamma \models_{\mathcal{A}} t \equiv_{\varepsilon} s$, if for any assignment $\iota: X \rightarrow A$, the following implication holds

$$(\forall t' \equiv_{\varepsilon'} s' \in \Gamma, d_A(\iota(t'), \iota(s')) \leq \varepsilon') \Rightarrow d_A(\iota(t), \iota(s)) \leq \varepsilon,$$

where $\iota(t)$ is the homomorphic interpretation of t in \mathcal{A} .

A quantitative algebra \mathcal{A} is said to *satisfy* (or be a *model* for) the quantitative theory \mathcal{U} , if $\Gamma \models_{\mathcal{A}} t \equiv_{\varepsilon} s$ whenever $\Gamma \vdash t \equiv_{\varepsilon} s \in \mathcal{U}$. We write $\mathbb{K}(\Sigma, \mathcal{U})$ for the collection of models of a theory \mathcal{U} of type Σ .

Sometimes it is convenient to consider the quantitative Σ -algebras whose carrier is a complete extended metric space. This class of algebras forms a full subcategory of $\mathbf{QA}(\Sigma)$, written $\mathbf{CQA}(\Sigma)$. Similarly, we write $\mathbb{CK}(\Sigma, \mathcal{U})$ for the full subcategory of quantitative Σ -algebras in $\mathbf{CQA}(\Sigma)$ which are models of \mathcal{U} .

The following lifts the Cauchy completion of metric spaces to quantitative algebras.

► **Definition 2.** (*Algebra Completion*) The Cauchy completion of a quantitative Σ -algebra $\mathcal{A} = (A, \Sigma^{\mathcal{A}})$, is the quantitative Σ -algebra $\overline{\mathcal{A}} = (\overline{A}, \Sigma^{\overline{\mathcal{A}}})$, where \overline{A} is the Cauchy completion of A and $\Sigma^{\overline{\mathcal{A}}} = \{f^{\overline{\mathcal{A}}}: \overline{A}^n \rightarrow \overline{A} \mid f: n \in \Sigma\}$ is such that for Cauchy sequences $(b_j^i)_j$ converging to $b^i \in \overline{A}$, for $1 \leq i \leq n$, $f^{\overline{\mathcal{A}}}(b^1, \dots, b^n) = \lim_j f^{\mathcal{A}}(b_j^1, \dots, b_j^n)$.

The above extends to a functor $\mathbb{C}: \mathbf{QA}(\Sigma) \rightarrow \mathbf{CQA}(\Sigma)$ which is the left adjoint to the functor embedding $\mathbf{CQA}(\Sigma)$ into $\mathbf{QA}(\Sigma)$.

The completion of quantitative Σ -algebras extends also to a functor from $\mathbb{K}(\Sigma, \mathcal{U})$ to $\mathbb{CK}(\Sigma, \mathcal{U})$, whenever \mathcal{U} can be axiomatised by a collection of *continuous schemata*, which are conditional quantitative equations of the form

$$\{x_i \equiv_{\varepsilon_i} y_i \mid i = 1..n\} \vdash t \equiv_{\varepsilon} s, \quad \text{for all } \varepsilon \geq f(\varepsilon_1, \dots, \varepsilon_n),$$

where $f: \mathbb{R}_{\geq 0}^n \rightarrow \mathbb{R}_{\geq 0}$ is a continuous real-valued function, $\varepsilon, \varepsilon_i \in \mathbb{Q}_{\geq 0}$, and $x_i, y_i \in X$. We call such a theory *continuous*.

Free Monads on Quantitative Equational Theories

To every signature Σ , one can associate a *signature endofunctor* (also called Σ) on \mathbf{Met} by:

$$\Sigma = \coprod_{f: n \in \Sigma} Id^n.$$

It is easy to see that, by couniversality of the coproduct, quantitative Σ -algebras correspond to Σ -algebras for the functor Σ in \mathbf{Met} , and the morphisms between them to non-expansive homomorphisms of Σ -algebras. Below we pass between the two points of view as convenient.

► **Theorem 3** (Free Algebra [21]). *The forgetful functor $\mathbb{K}(\Sigma, \mathcal{U}) \rightarrow \mathbf{Met}$ has a left adjoint.*

The left adjoint assigns to any $X \in \mathbf{Met}$ a *free quantitative Σ -algebra* $(T_X, \psi_X^{\mathcal{U}})$ satisfying \mathcal{U} , from which one canonically obtains the monad $(T_{\mathcal{U}}, \eta^{\mathcal{U}}, \mu^{\mathcal{U}})$, with functor $T_{\mathcal{U}}: \mathbf{Met} \rightarrow \mathbf{Met}$ mapping $X \in \mathbf{Met}$ to the carrier T_X of the free algebra.

A similar free construction also holds for quantitative algebras in $\mathbf{CQA}(\Sigma)$ for continuous quantitative equational theories, implying that the forgetful functor from $\mathbb{CK}(\Sigma, \mathcal{U})$ to \mathbf{CMet} has a left adjoint. In particular, $\mathbb{C}T_{\mathcal{U}}$ is the free monad on \mathcal{U} in \mathbf{CMet} , provided that the quantitative equational theory is continuous.

Finally, let $T\text{-Alg}$ denote the category of Eilenberg-Moore (EM) algebras for a monad T . In [1], it is shown that, whenever the quantitative theory \mathcal{U} is *basic*, *i.e.*, it can be axiomatised by a set of conditional equations of the form

$$\{x_1 \equiv_{\varepsilon_1} y_1, \dots, x_n \equiv_{\varepsilon_n} y_n\} \vdash t \equiv_{\varepsilon} s,$$

where $x_i, y_i \in X$ (*cf.* [22]), then EM $T_{\mathcal{U}}$ -algebras are in 1-1 correspondence with the quantitative algebras satisfying \mathcal{U} :

► **Theorem 4.** *For any basic quantitative equational theory \mathcal{U} of type Σ , $T_{\mathcal{U}}\text{-Alg} \cong \mathbb{K}(\Sigma, \mathcal{U})$.*

4

 Tensor of Strong Monads

In this section we provide the definition of *tensor of strong monads* on a generic symmetric monoidal closed category \mathbf{V} . The presentation follows and generalises that of Manes [20], which considers only the case of monads on \mathbf{Set} .

Let v be an object in \mathbf{V} . As \mathbf{V} is self-enriched, it has all v -fold *powers* (or v -*powers*) X^v , of any object $X \in \mathbf{V}$, defined as $X^v = [v, X]$ [16]. Moreover, $(-)^v: \mathbf{V} \rightarrow \mathbf{V}$ is a strong functor with strength $\xi_{X,Y}: X \square Y^v \rightarrow (X \square Y)^v$ obtained by currying

$$v \square (X \square Y^v) \xrightarrow{\cong} X \square (v \square Y^v) \xrightarrow{X \square ev} X \square Y.$$

Let $F: \mathbf{V} \rightarrow \mathbf{V}$ be a strong functor with strength t . The v -*power* functor $(-)^v$ can be lifted to F -algebras by mapping (A, a) to $(A, a)^v = (A^v, a^v \circ \sigma_A)$, where $\sigma_A: FA^v \Rightarrow (FA)^v$ is the strong natural transformation obtained from t by currying $Fev_A^v \circ t_{v, A^v}$. Hence F -algebras are closed under powers of \mathbf{V} -objects.

► **Definition 5** (Pre-operation of a strong functor). *Let $F: \mathbf{V} \rightarrow \mathbf{V}$ be a strong functor and $v \in \mathbf{V}$. A v -ary pre-operation of F is a strong natural transformation of type $(-)^v \Rightarrow F$.*

We denote by $\mathcal{O}_F(v)$ the set of v -ary pre-operations of F . An application of $g \in \mathcal{O}_F(v)$ to an F -algebra (A, a) is the composite $a^g = a \circ g_A$. We call a^g an *operation* of (A, a) .

► **Proposition 6.** *Let $(A, a), (B, b)$ be F -algebras of a strong endofunctor F on \mathbf{V} and $f: A \rightarrow B$ a morphism in \mathbf{V} . Then, the following are equivalent:*

1. f is an F -homomorphism from (A, a) to (B, b) ;
2. For every $v \in \mathbf{V}$ and $g \in \mathcal{O}_F(v)$, $f \circ a^g = b^g \circ f^v$.

The above proposition indicates that F -algebras are precisely characterised by their operations. In some situations, depending on the functor F , one gets the same characterisation with much fewer operations. We identify this property with the following definition.

► **Definition 7** (Density). *A set \mathcal{D} of pre-operations of a strong functor $F: \mathbf{V} \rightarrow \mathbf{V}$ is dense, if for any F -algebras $(A, a), (B, b)$ and $f: A \rightarrow B$ in \mathbf{V} , the following are equivalent:*

1. f is an F -homomorphism from (A, a) to (B, b) ;
2. For every v -ary pre-operation $g \in \mathcal{D}$, $f \circ a^g = b^g \circ f^v$.

Let F, G be two strong endofunctors on \mathbf{V} . A $\langle F, G \rangle$ -*bialgebra* is a triple (A, a, b) consisting of an object $A \in \mathbf{V}$ with both an F -algebra structure $a: FA \rightarrow A$ and a G -algebra structure $b: GA \rightarrow A$. A morphism of $\langle F, G \rangle$ -bialgebras is an arrow that is simultaneously an F - and G -homomorphism. Denote by $\langle F, G \rangle$ -**biAlg** the category of $\langle F, G \rangle$ -bialgebras.

► **Proposition 8.** *Let (A, a, b) be an $\langle F, G \rangle$ -bialgebra. The following statements are equivalent:*

1. For all $v \in \mathbf{V}$ and $g \in \mathcal{O}_F(v)$, a^g is a G -homomorphism;
2. For all $w \in \mathbf{V}$ and $h \in \mathcal{O}_G(w)$, b^h is an F -homomorphism.

Diagrammatically:

$$\begin{array}{ccc} GA^v \xrightarrow{\bar{b}} A^v & & FA^w \xrightarrow{\bar{a}} A^w \\ \downarrow^{G(a^g)} \quad (1) \quad \downarrow^{a^g} & \text{iff} & \downarrow^{F(b^h)} \quad (2) \quad \downarrow^{b^h} \\ GA \xrightarrow{b} A & & FA \xrightarrow{a} A \end{array}$$

where $(A, a)^w = (A^w, \bar{a})$ and $(A, b)^v = (A^v, \bar{b})$.

► **Definition 9** (Commutative bialgebra). A $\langle F, G \rangle$ -bialgebra (A, a, b) is commutative if it satisfies either of the equivalent conditions of Proposition 8.

In the case the functors F and G admit dense sets of pre-operations, commutativity for their bialgebras can be more conveniently expressed in the following way.

► **Proposition 10.** Let \mathcal{D} and \mathcal{E} be dense sets of pre-operations for F and G , respectively. A $\langle F, G \rangle$ -bialgebra (A, a, b) is commutative iff it satisfies either of the equivalent conditions:

1. For all $g \in \mathcal{D}$, a^g is a G -homomorphism;
2. For all $h \in \mathcal{E}$, b^h is an F -homomorphism.

Let (T, η, μ) be a strong monad on \mathbf{V} . Note that, as T is a strong functor and the EM-algebras for T are closed under powers of \mathbf{V} -objects, all the results and definitions given in this section extends to EM-algebras for T .

Let (T, η, μ) , (T', η', μ') be two strong monads on \mathbf{V} . A EM $\langle T, T' \rangle$ -bialgebra is a triple (A, a, a') consisting of an object $A \in \mathbf{V}$ with both an EM T -algebra structure $a: TA \rightarrow A$ and an EM T' -algebra structure $a': T'A \rightarrow A$. We say that an EM $\langle T, T' \rangle$ -bialgebra (A, a, a') is *commutative* if it is so as a $\langle T, T' \rangle$ -bialgebra for the functors T, T' . We denote by $\langle T, T' \rangle$ -**biAlg** the category of EM $\langle T, T' \rangle$ -bialgebras and by $(T \otimes T')$ -**biAlg**, the full subcategory of the commutative EM $\langle T, T' \rangle$ -bialgebras.

► **Definition 11** (Tensor of monads). If the forgetful functor $(T \otimes T')$ -**biAlg** $\rightarrow \mathbf{V}$ has a left adjoint, then the monad induced by the adjunction is the tensor of T, T' , denoted $T \otimes T'$.

Note that the tensor of monads does not necessarily exist (see [4] for counterexamples). However, when it does $T \otimes T' \cong T' \otimes T$, as the categories of commutative biagebras $(T \otimes T')$ -**biAlg** and $(T' \otimes T)$ -**biAlg** are isomorphic.

5 Tensor of Quantitative Theories

In this section, we develop the theory for the *tensor* of quantitative equational theories. The main result is that the free monad on the tensor of two theories is the tensor of the monads on the theories. In the proof given, we use the fact that the quantitative theories are *basic*, as this allows us to exploit the correspondence between the algebras of a theory \mathcal{U} and the EM-algebras of the monad $T_{\mathcal{U}}$ (Theorem 4).

Let Σ, Σ' be two disjoint signatures. Following Freyd [8] (and [12]), we define the tensor of two quantitative equational theories $\mathcal{U}, \mathcal{U}'$ of respective types Σ and Σ' , written $\mathcal{U} \otimes \mathcal{U}'$, as the smallest quantitative theory containing $\mathcal{U}, \mathcal{U}'$ and the quantitative equations

$$\vdash f(g(x_1^1, \dots, x_m^1), \dots, g(x_1^n, \dots, x_m^n)) \equiv_0 g(f(x_1^1, \dots, x_1^n), \dots, f(x_m^1, \dots, x_m^n)), \quad (1)$$

for all $f: n \in \Sigma$ and $g: m \in \Sigma'$, expressing that the operations of one theory commute with the operations of the other.

5.1 Density of Symbolic Pre-operations

Towards our main result, we identify a dense set of pre-operations for the free monads on quantitative equational theories which, in turn, will gives us a simpler characterization for commutative bialgebras for these monads (*cf.* Proposition 10).

First notice that any signature functor $\Sigma = \coprod_{f:n \in \Sigma} Id^n$ in **Met** is strong, as it is the coproduct of the strong functors $Id^n \cong (-)^{\underline{n}}$, where $\underline{n} \in \mathbf{Met}$ denotes the set $\{1, \dots, n\}$

7:8 Tensor of Quantitative Equational Theories

equipped with the discrete extended metric assigning infinite distance to distinct elements. Moreover, the injections $in_f: (-)^n \Rightarrow \Sigma$ are strong natural transformations, hence they are n -ary pre-operations of Σ (cf. Definition 5).

► **Proposition 12.** $\mathcal{S}_\Sigma = \{in_f \mid f: n \in \Sigma\}$ is a dense set of pre-operations of Σ .

In the following, the pre-operations in \mathcal{S}_Σ will be called *symbolic*, and to simplify the notation, for any $f: n \in \Sigma$ and Σ -algebra (A, a) , we write a^f instead of a^{in_f} .

Let \mathcal{U} be a quantitative equational theory of type Σ . Then, also the monad $T_{\mathcal{U}}$ is strong, with strength $\zeta_{X,Y}: X \square T_{\mathcal{U}}Y \rightarrow T_{\mathcal{U}}(X \square Y)$ obtained by uncurrying the unique map $h_{X,Y}$ that, by Theorem 3, makes the following diagram commute

$$\begin{array}{ccc} Y & \xrightarrow{\eta_Y^{\mathcal{U}}} & T_{\mathcal{U}}Y & \xleftarrow{\psi_Y^{\mathcal{U}}} & \Sigma T_{\mathcal{U}}Y \\ \beta_{X,Y} \searrow & & \downarrow h_{X,Y} & & \downarrow \Sigma h_{X,Y} \\ & & (T_{\mathcal{U}}(X \square Y))^X & \xleftarrow{\psi_{(T(X \square Y))^X}^{\mathcal{U}}} & \Sigma(T_{\mathcal{U}}(X \square Y))^X \end{array}$$

where $\beta_{X,Y}$ is the currying of $\eta_{X \square Y}^{\mathcal{U}}: X \square Y \rightarrow T_{\mathcal{U}}(X \square Y)$.

Since a monad is strong iff both its unit and multiplication are strong natural transformations, both $\eta^{\mathcal{U}}$, $\mu^{\mathcal{U}}$ are strong. Moreover, also $\psi^{\mathcal{U}}: \Sigma T_{\mathcal{U}} \Rightarrow T_{\mathcal{U}}$ is strong.

Thus any pre-operation $g \in \mathcal{O}_\Sigma(v)$ can be tuned into an pre-operation of $T_{\mathcal{U}}$ as the composite

$$(-)^v \xrightarrow{g} \Sigma \xrightarrow{\Sigma \eta^{\mathcal{U}}} \Sigma T_{\mathcal{U}} \xrightarrow{\psi^{\mathcal{U}}} T_{\mathcal{U}}.$$

In particular, when the theory \mathcal{U} is basic, by exploiting Theorem 4, the above transformation allows us to turn any dense set of pre-operations of Σ into a dense set of pre-operations of $T_{\mathcal{U}}$.

► **Proposition 13.** Let \mathcal{U} be a basic quantitative theory of type Σ and \mathcal{D} a dense set of pre-operations of Σ . Then $\{\psi^{\mathcal{U}} \circ \Sigma \eta^{\mathcal{U}} \circ g \mid g \in \mathcal{D}\}$ is a dense set of pre-operations of $T_{\mathcal{U}}$.

By combining Propositions 12 and 13, we have that $\mathcal{S}_{T_{\mathcal{U}}} = \{\psi^{\mathcal{U}} \circ \Sigma \eta^{\mathcal{U}} \circ in_f \mid f: n \in \Sigma\}$ is a dense set of pre-operations of $T_{\mathcal{U}}$. We call also these pre-operations *symbolic* and we simplify the notation by writing $a^{(f)}$ instead of $a^{(\psi^{\mathcal{U}} \circ \Sigma \eta^{\mathcal{U}} \circ in_f)}$, for $f: n \in \Sigma$ and $(A, a) \in T_{\mathcal{U}}\text{-Alg}$.

Thus, as an immediate consequence of Propositions 10 and 13, we obtain the following simpler characterization for commutative $\langle T_{\mathcal{U}}, T_{\mathcal{U}'} \rangle$ -bialgebras.

► **Corollary 14.** Let $\mathcal{U}, \mathcal{U}'$ be basic quantitative theories respectively of type Σ, Σ' . A $\langle T_{\mathcal{U}}, T_{\mathcal{U}'} \rangle$ -bialgebra (A, a, b) is commutative iff it satisfies either of the equivalent conditions

1. For all $f: n \in \Sigma$, $a^{(f)}$ is a $T_{\mathcal{U}'}$ -homomorphism;
2. For all $g: m \in \Sigma'$, $b^{(g)}$ is a $T_{\mathcal{U}}$ -homomorphism.

5.2 Tensor of Free Monads on Quantitative Theories

Let $\mathcal{U}, \mathcal{U}'$ be basic quantitative theories respectively of type Σ, Σ' . We show that any model for $\mathcal{U} \otimes \mathcal{U}'$ is a $\langle \mathcal{U} \otimes \mathcal{U}' \rangle$ -bialgebra: an extended metric space A with both a Σ -algebra structure $a: \Sigma A \rightarrow A$ satisfying \mathcal{U} and a Σ' -algebra structure $b: \Sigma' A \rightarrow A$ satisfying \mathcal{U}' and respecting the diagrammatic condition below, for all $f: n \in \Sigma$ and $g: m \in \Sigma'$

$$\begin{array}{ccc} A^n & \xrightarrow{a^f} & A & \xleftarrow{b^g} & A^m \\ (b^g)^n \uparrow & & & & \uparrow (a^f)^m \\ (A^m)^n & \xrightarrow[\cong]{\chi} & & & (A^n)^m \end{array} \quad (2)$$

Formally, we denote by $(\mathcal{U} \otimes \mathcal{U}')$ -**biAlg** the category of $\langle \mathcal{U} \otimes \mathcal{U}' \rangle$ -bialgebras, with morphisms the non-expansive homomorphisms preserving both algebraic structures. Then, the following isomorphism of categories holds.

► **Proposition 15.** $\mathbb{K}(\Sigma + \Sigma', \mathcal{U} \otimes \mathcal{U}') \cong (\mathcal{U} \otimes \mathcal{U}')\text{-biAlg}$, for $\mathcal{U}, \mathcal{U}'$ basic quantitative theories.

Moreover, by adapting the isomorphism of Theorem 4 and exploiting the density of symbolic pre-operations (cf. Corollary 14) the following is also true.

► **Proposition 16.** $(\mathcal{U} \otimes \mathcal{U}')\text{-biAlg} \cong (T_{\mathcal{U}} \otimes T_{\mathcal{U}'})\text{-biAlg}$, for $\mathcal{U}, \mathcal{U}'$ basic quantitative theories.

By combining the above two propositions we get the main theorem of this section.

► **Theorem 17.** Let $\mathcal{U}, \mathcal{U}'$ be basic quantitative theories. Then, the monad $T_{\mathcal{U} \otimes \mathcal{U}'}$ in **Met** is the tensor of monads $T_{\mathcal{U}} \otimes T_{\mathcal{U}'}$.

Proof. By Propositions 15 and 16 the forgetful functor from $(T_{\mathcal{U}} \otimes T_{\mathcal{U}'})\text{-biAlg}$ to **Met** has a left adjoint and the monad generated by this adjunction is isomorphic to $T_{\mathcal{U} \otimes \mathcal{U}'}$. Thus, by definition of tensor of monads, $T_{\mathcal{U} \otimes \mathcal{U}'} \cong T_{\mathcal{U}} \otimes T_{\mathcal{U}'}$. ◀

The above results do not require any specific property of **Met**, apart that its morphisms are non-expansive maps. Thus, when the quantitative equational theories are continuous, we can reformulate an alternative version of Theorem 17 which is valid in **CMet**.

► **Theorem 18.** Let $\mathcal{U}, \mathcal{U}'$ be continuous quantitative theories. Then, $\mathbb{C}T_{\mathcal{U} \otimes \mathcal{U}'}$ in **CMet** is the tensor of monads $\mathbb{C}T_{\mathcal{U}} \otimes \mathbb{C}T_{\mathcal{U}'}$.

6 Quantitative Reader Algebras

Let E be a finite set of input values and fix an enumeration $E = \{e_1, \dots, e_n\}$ for it. The *quantitative reader algebras* of type E are the algebras for the signature

$$\Sigma_{\mathcal{R}_E} = \{r: |E|\}$$

having only one operator r of arity equal to the number of the input values in E , and satisfying the following axioms

$$(\text{Idem}) \vdash x \equiv_0 r(x, \dots, x),$$

$$(\text{Diag}) \vdash r(x_{1,1}, \dots, x_{n,n}) \equiv_0 r(r(x_{1,1}, \dots, x_{1,n}), \dots, r(x_{n,1}, \dots, x_{n,n})).$$

The quantitative theory induced by the axioms above, written \mathcal{R}_E , is called *quantitative theory of reading computations*.

Intuitively, the term $r(t_1, \dots, t_n)$ can be interpreted as the computation that proceeds as t_i after reading the value e_i from its input. The axiom (Idem) says that if we ignore the value of the input the reading of it is not observable; (Diag) says that the resulting computation after reading the input is the same no matter how many times we read it.

► **Remark 19.** For the binary case ($|E| = 2$) we can think of r as an *if-then-else* statement $b?(S, T)$ checking for the value of a fixed global Boolean variable b and proceeding as S when $b = \text{true}$, and as T otherwise. In this case, (Idem) and (Diag) express the standard program equivalences $S \equiv b?(S, S)$ and $b?(S, T) \equiv b?(b?(S, T'), b?(S', T))$.

In the following, when the set E is clear from the context, we use \mathcal{R} in place of \mathcal{R}_E .

On Metric Spaces

Let E be a finite set. We denote by \underline{E} the extended metric space on E equipped with the indiscrete metric that assigns infinite distance to any pair of distinct elements.

Consider the \underline{E} -power functor $(-)^{\underline{E}}: \mathbf{Met} \rightarrow \mathbf{Met}$, assigning to each $X \in \mathbf{Met}$ the metric space $[E, X]$ of (necessarily non-expansive) maps from \underline{E} to X .

This functor has a monad structure, with unit $\kappa: Id \Rightarrow (-)^{\underline{E}}$ and multiplication $\Delta: ((-)^{\underline{E}})^{\underline{E}} \Rightarrow (-)^{\underline{E}}$, respectively given as follows, for $x \in X$, $e \in E$, and $f \in E \rightarrow X^{\underline{E}}$

$$\kappa_X(x)(e) = x, \quad \Delta_X(f)(e) = f(e)(e).$$

This is also known as *reader monad* (also called *environment monad* or *function monad*).

► **Remark 20.** The reader monad is always well defined in a cartesian closed category. Fix an object E . The reader monad $(-)^E$ has unit and multiplication respectively given by

$$X \cong X^1 \xrightarrow{X^!} X^E \quad \text{and} \quad (X^E)^E \cong X^{E \times E} \xrightarrow{X^\delta} X^E,$$

where $!: E \rightarrow 1$ is the unique map to the terminal object and $\delta: E \rightarrow E \times E$ the diagonal map $\delta = \langle id, id \rangle$. However, this definition does not generalise to arbitrary monoidal closed categories, and \mathbf{Met} is such a counterexample. The specific problem with \mathbf{Met} is that $\delta: E \rightarrow E \square E$ is not well-defined for arbitrary $E \in \mathbf{Met}$, as non-expansiveness requires that

$$d_E(e, e') \geq d_{E \square E}(\delta(e), \delta(e')) = d_E(e, e') + d_E(e, e'),$$

which holds only when E has the discrete metric. This is the reason why in our treatment we restrict the set of input values to have discrete metric.

The reader monad $(-)^{\underline{E}}$ is isomorphic to the free monad $T_{\mathcal{R}}$. In other words, the quantitative theory \mathcal{R} of reading computations axiomatises the reader monad.

► **Theorem 21.** *The monads $T_{\mathcal{R}}$ and $(-)^{\underline{E}}$ in \mathbf{Met} are isomorphic.*

Let T be a strong monad with strength t . The natural transformation $\lambda_X: TX^{\underline{E}} \Rightarrow (TX)^{\underline{E}}$ obtained from the strength t by currying $Tev_X^{\underline{E}} \circ t_{\underline{E}, X^{\underline{E}}}$, is a distributive law of monads. Distributive laws induce a notion of monad composition [2], so Moggi's reader monad transformer $T \mapsto (T-)^{\underline{E}}$ is also available in \mathbf{Met} . The following says that we can recover this monad transformer as the operation of tensoring with the reader monad.

► **Theorem 22 (Tensoring with Reader Monad).** *Let T be a strong monad. Then, $T \otimes (-)^{\underline{E}}$ exists and is given as the monad composition $(T-)^{\underline{E}}$.*

By using the above result in combination with Theorem 17, we obtain an analogous transformer at the level of quantitative equational theories as follows.

► **Corollary 23.** *Let \mathcal{U} be a basic quantitative equational theory. Then, $(T_{\mathcal{U}}-)^{\underline{E}}$ is the free monad on the theory $\mathcal{U} \otimes \mathcal{R}$ in \mathbf{Met} .*

On Complete Metric Spaces

The category \mathbf{CMet} has finite products. Since, we assumed the set of input values E to be finite, the functor $(-)^{\underline{E}}$ is isomorphic to the finite product $(-)^n$, for $n = |E|$. Therefore the power functor $(-)^{\underline{E}}$, preserves Cauchy completeness and can be restricted to an endofunctor on \mathbf{CMet} . Thus also the reader monad restricts to \mathbf{CMet} .

Because \mathcal{R} is a continuous quantitative theory, the free monad on \mathcal{R} in \mathbf{CMet} is $\mathcal{CT}_{\mathcal{R}}$. Thus, by restricting Theorem 21 to quantitative algebras over \mathbf{CMet} , we obtain:

► **Theorem 24.** *The monads $\mathbb{C}T_{\mathcal{R}}$ and $(-)^{\underline{E}}$ in \mathbf{CMet} are isomorphic.*

In virtue of the above characterisation, by instantiating Theorem 22 in the category of complete extended metric spaces, in combination with Theorems 17 we obtain the following variant of the quantitative reader theory transformer on continuous quantitative theories.

► **Corollary 25.** *Let \mathcal{U} be a continuous quantitative theory. Then, $(\mathbb{C}T_{\mathcal{U}}-)^{\underline{E}}$ is the free monad on the theory $\mathcal{U} \otimes \mathcal{R}$ in \mathbf{CMet} .*

7 Quantitative Writer Algebras

Fix an extended metric space $\Lambda \in \mathbf{Met}$ of *output values* having monoid structure $(\Lambda, *, 0)$ with non-expansive multiplication operation $*$: $\Lambda \times \Lambda \rightarrow \Lambda$.

The *quantitative writer algebras* of type Λ are the algebras for the signature

$$\Sigma_{\mathcal{W}_{\Lambda}} = \{\mathbf{w}_{\alpha} : 1 \mid \alpha \in \Lambda\}$$

having a unary operator \mathbf{w}_{α} , for each output value $\alpha \in \Lambda$, and satisfying the following axioms

$$\text{(Zero)} \vdash x \equiv_0 \mathbf{w}_0(x),$$

$$\text{(Mult)} \vdash \mathbf{w}_{\alpha}(\mathbf{w}_{\alpha'}(x)) \equiv_0 \mathbf{w}_{\alpha * \alpha'}(x),$$

$$\text{(Diff)} \{x \equiv_{\varepsilon} x'\} \vdash \mathbf{w}_{\alpha}(x) \equiv_{\delta} \mathbf{w}_{\alpha'}(x'), \text{ for } \delta \geq d_{\Lambda}(\alpha, \alpha') + \varepsilon.$$

The quantitative theory induced by the axioms above, written \mathcal{W}_{Λ} , is called *quantitative theory of writing computations*.

The term $\mathbf{w}_{\alpha}(t)$ represents the computation that proceeds as t after writing α on the output tape. The axiom (Zero) says that writing the identity element 0 is not observable on the tape; (Mult) says that consecutive writing operations are stored in the tape in the order of execution; (Diff) compares two computations w.r.t. the distance of their output values.

In the following, when the metric space Λ of output values is clear from the context, we use \mathcal{W} in place of \mathcal{W}_{Λ} .

On Metric Spaces

Let $(\Lambda \square -): \mathbf{Met} \rightarrow \mathbf{Met}$ be the functor assigning to each extended metric space X the space $(\Lambda \square X)$. By exploiting the monoid structure of Λ , the functor $(\Lambda \square -)$ can be given a monad structure with unit $\tau: Id \Rightarrow (\Lambda \square -)$ and multiplication $\varsigma: (\Lambda \square (\Lambda \square -)) \Rightarrow (\Lambda \square -)$, respectively given as follows, for arbitrary $x \in X$ and $\alpha, \alpha' \in \Lambda$

$$\tau_X(x) = (0, x),$$

$$\varsigma_X((\alpha, (\alpha', x))) = (\alpha * \alpha', x).$$

This monad is also known as *writer monad* (also called *complexity monad*). Note that, the non-expansiveness of the maps above crucially depends on the assumption that the multiplication $*$ in Λ is non-expansive.

The next theorem says that the writer monad $(\Lambda \square -)$ has a quantitative equational presentation in terms of the theory \mathcal{W} of writing computations.

► **Theorem 26.** *The monads $T_{\mathcal{W}}$ and $(\Lambda \square -)$ in \mathbf{Met} are isomorphic.*

7:12 Tensor of Quantitative Equational Theories

Let T be a strong monad with strength t . There is a canonical distributive law of the monad $(\Lambda \square -)$ over T , obtained using the strength $t_{\Lambda, -} : (\Lambda \square T-) \Rightarrow T(\Lambda \square -)$ of T . So $T(\Lambda \square -)$ acquires a canonical monad structure [2], and we obtain Moggi's writer monad transformer $T \mapsto T(\Lambda \square -)$ in **Met**.

In [12], Hyland et al. observed that Moggi's writer monad transformer can be equivalently recovered as the operation of tensoring with the writer monad.

► **Theorem 27** (Tensoring with Writer Monad [12]). *Let T be a strong monad. Then, the monad composition $T(\Lambda \square -)$ is given as $T \otimes (\Lambda \square -)$.*

By combining the above with Theorems 17 and 26, we get an analogous transformer at the level of quantitative equational theories as follows:

► **Corollary 28.** *Let \mathcal{U} be a basic quantitative theory. Then, $T_{\mathcal{U}}(\Lambda \square -)$ is the free monad on the theory $\mathcal{U} \otimes \mathcal{W}$ in **Met**.*

On Complete Metric Spaces

If we assume the monoid $(\Lambda, *, 0)$ to be over a complete extended metric space Λ , the writer monad $(\Lambda \square -)$ is well defined also in **CMet**.

Since \mathcal{W} is axiomatised by a continuous schema of quantitative conditional equations the free monad on \mathcal{W} in **CMet** is given by $\mathcal{CT}_{\mathcal{W}}$. Thus, by restricting the use of Theorem 26 to quantitative algebras over complete extended metric spaces, we obtain:

► **Theorem 29.** *The monads $\mathcal{CT}_{\mathcal{W}}$ and $(\Lambda \square -)$ in **CMet** are isomorphic.*

Thus, by similar arguments as before, we obtain the following variant of Corollary 28.

► **Corollary 30.** *Let \mathcal{U} be a continuous quantitative theory. Then, $\mathcal{CT}_{\mathcal{U}}(\Lambda \square -)$ is the free monad on the theory $\mathcal{U} \otimes \mathcal{W}$ in **CMet**.*

8 The Algebras of Labeled Markov Processes

In this section we show how to obtain a quantitative equational axiomatization of labelled Markov processes with discounted bisimilarity metric as the combination, via sum and tensor, of the following simpler quantitative equational theories:

- (a) *The quantitative theory of interpolative barycentric algebras \mathcal{B} from [21] over the signature $\Sigma_{\mathcal{B}} = \{+_e : 2 \mid e \in [0, 1]\}$ extends M. H. Stone's theory of barycentric algebras [31] (a.k.a. abstract convex algebras) with the following axiom*

$$(IB) \quad \{x \equiv_{\varepsilon} y, x' \equiv_{\varepsilon'} y'\} \vdash x +_e x' \equiv_{\delta} y +_e y', \text{ for } \delta \geq e\varepsilon + (1 - e)\varepsilon'$$

expressing that the distance between convex combinations is obtained as the convex interpolation of the distance of their sub-terms. This theory will be used to axiomatise probability distributions with Kantorovich metric [15].

- (b) *The pointed quantitative theory, defined as the free quantitative theory $\mathcal{U}_{\mathbf{0}}$ (i.e., the one imposing no additional axioms) for a signature $\Sigma_{\mathbf{0}} = \{\mathbf{0} : 0\}$ consisting of a single constant $\mathbf{0}$ symbol. This will be used to axiomatise termination.*
- (c) *The quantitative theory \mathcal{R}_A of reading computations (cf. Section 6) will be used to axiomatise the reaction to the choice of a label from a set A of action labels.*

- (d) *The quantitative theory of contractive operators* discussed in [1], is the theory obtained by imposing a Lipschitz contractive axiom for each operator in the signature. In our case, we consider a signature $\Sigma_\diamond = \{\diamond: 1\}$ with only one unary operator and the contractive theory \mathcal{U}_\diamond generated from the axiom

$$(\diamond\text{-Lip}) \quad \{x =_\varepsilon y\} \vdash \diamond(x) \equiv_\delta \diamond(y), \text{ for } \delta \geq c\varepsilon,$$

where $c \in (0, 1)$ is a fixed *contractive factor* for the operator \diamond . This theory will be used to axiomatise the transition to a next state with discount factor c .

Formally, we define the quantitative theory \mathcal{U}_{LMP} of labelled Markov processes as the following combination of quantitative theories, with signature Σ_{LMP} given by the disjoint union of those from its component theories:

$$\Sigma_{\text{LMP}} = \Sigma_{\mathcal{B}} + \Sigma_{\mathbf{0}} + \Sigma_{\mathcal{R}_A} + \Sigma_\diamond, \quad \mathcal{U}_{\text{LMP}} = ((\mathcal{B} + \mathcal{U}_{\mathbf{0}}) \otimes \mathcal{R}_A) + \mathcal{U}_\diamond.$$

Following [32, Section 6], we regard A -labelled Markov processes over extended metric spaces as $(\Delta(1 + -))^{\mathbb{A}}$ -coalgebras in \mathbf{Met} , where Δ is the *Kantorovich functor* assigning to each $X \in \mathbf{Met}$ the space of Radon probability measures with finite moment over X equipped with Kantorovich metric. In [32] it is shown that the *probabilistic bisimilarity distance* on a labelled Markov processes (X, τ) is equal to the (pseudo)metric

$$\mathbf{d}_{(X, \tau)}(x, x') = d_Z(h(x), h(x')),$$

where $h: X \rightarrow Z$ is the unique homomorphism to the final coalgebra (Z, ω) .

Similarly to [1], we slightly extend the type of the coalgebras to encompass the case when the probabilistic bisimilarity distance is discounted by a factor $0 < c < 1$. Explicitly, we consider coalgebras for the functor $(\Delta(1 + c \cdot -))^{\mathbb{A}}$, where $(c \cdot -)$ is the *rescaling functor*, mapping a metric space (X, d_X) to $(X, c \cdot d_X)$. This will not change the essence of the results from [32] that are used in this section to characterise the probabilistic bisimilarity metric.

In the remainder of the section we prove that the theory \mathcal{U}_{LMP} axiomatizes (the monad of) A -labelled Markov processes with c -discounted bisimilarity metric.

On Metric Spaces

We characterise the monad $T_{\mathcal{U}_{\text{LMP}}}$ in steps. First, note that $T_{\mathcal{U}_{\mathbf{0}}} \cong 1^* = (1 + -)$ is the *maybe monad*, i.e., freely generated monad on the constant terminal object functor 1 . As the monad $(1 + -)$ is isomorphic to $(1F)^*$, for any functor F , by [1, Theorems 4.4 and 5.2], and [12, Theorem 4], we obtain the following isomorphism of monads in \mathbf{Met} :

$$T_{\mathcal{B} + \mathcal{U}_{\mathbf{0}}} \cong T_{\mathcal{B}} + T_{\mathcal{U}_{\mathbf{0}}} \cong \Pi(1 + -),$$

where $\Pi(1 + -)$ is the *finite sub-distribution monad* with functor assigning to $X \in \mathbf{Met}$ the space of finitely supported Borel sub-probability measures with Kantorovich metric. Thus, $\mathcal{B} + \mathcal{U}_{\mathbf{0}}$ axiomatizes finitely supported sub-probability distributions with Kantorovich metric.

From the above, Theorem 17 and Corollary 23, we further get the monad isomorphism

$$T_{(\mathcal{B} + \mathcal{U}_{\mathbf{0}}) \otimes \mathcal{R}_A} \cong \Pi(1 + -) \otimes (-)^{\mathbb{A}} \cong (\Pi(1 + -))^{\mathbb{A}},$$

saying that tensoring with the theory \mathcal{R}_A of reading computations corresponds to axiomatically adding the capability of reacting to the choice of an action label.

By [1, Theorem 6.3], $T_{\mathcal{U}_\diamond}$ is isomorphic to the free monad over the rescaling functor $(c \cdot -)$. Hence, by [1, Theorem 4.4] and [12, Corollary 2] we get the following last isomorphism

$$T_{\mathcal{U}_{\mathbf{LMP}}} = T_{((\mathcal{B} + \mathcal{U}_\diamond) \otimes \mathcal{R}_A) + \mathcal{U}_\diamond} \cong \mu y. (\Pi(1 + c \cdot y + -))^A.$$

Explicitly, this means that, the free monad on $\mathcal{U}_{\mathbf{LMP}}$ assigns to an arbitrary metric space $X \in \mathbf{Met}$ the *initial solution* of the following functorial equation in \mathbf{Met}

$$LMP_X \cong (\Pi(1 + c \cdot LMP_X + X))^A.$$

In particular, when $X = 0$ is the empty metric space (*i.e.*, the initial object in \mathbf{Met}) the above corresponds to the isomorphism on the initial $(\Pi(1 + c \cdot -))^A$ -algebra. The isomorphism gives us also a $(\Pi(1 + c \cdot -))^A$ -coalgebra structure on LMP_0 , which can be converted into a labeled Markov process (LMP_0, τ_0) via a post-composition with the inclusion $\Pi(-) \hookrightarrow \Delta(-)$.

The key aspect is that the metric of LMP_0 is exactly the bisimilarity metric.

► **Lemma 31.** d_{LMP_0} is the c -discounted probabilistic bisimilarity metric on (LMP_0, τ_0) .

► **Remark 32.** For a less abstract description of (LMP_0, τ_0) , notice that the elements of LMP_0 are (equivalence classes of) ground terms over the signature $\Sigma_{\mathbf{LMP}}$, which one can interpret as pointed (or rooted) acyclic labelled Markov processes quotiented by bisimilarity.

On Complete Metric Spaces

Since all the quantitative theories considered are continuous, we can replicate the same steps also while interpreting the theory $\mathcal{U}_{\mathbf{LMP}}$ over complete metric spaces, obtaining the monad

$$\mathbb{C}T_{\mathcal{U}_{\mathbf{LMP}}} \cong \mu y. \Delta(1 + c \cdot y + -)^A.$$

By following similar arguments to [1, Section 8.3], one can prove that the functorial equation $LMP_X \cong \Delta(1 + c \cdot LMP_X + X)^A$ has a unique solution. Thus by applying the monad above on $X = 0$ we recover the carrier of the final $(\Delta(1 + c \cdot -))^A$ -coalgebra, equipped with c -discounted probabilistic bisimilarity metric.

► **Remark 33.** While by interpreting the theory $\mathcal{U}_{\mathbf{LMP}}$ over \mathbf{Met} we can only characterise Markov processes that are acyclic, by doing it over \mathbf{CMet} we obtain an algebraic representation of all bisimilarity classes as the elements of the final coalgebra. Thus, among others, we also recover Markov processes with cyclic structures as the limit of all their finite unfoldings.

9 The Algebras of Markov Decision Processes with Rewards

As a last example, we provide a quantitative axiomatization of Markov decision processes with rewards equipped with discounted bisimilarity metric. As the construction is similar to Section 8, we avoid repeating the details of each step of the monad characterization.

Let $(\mathbb{R}, +, 0)$ be the standard monoid structure on the reals. We define the quantitative theory $\mathcal{U}_{\mathbf{MDP}}$ of Markov decision processes with real-valued rewards as follows

$$\Sigma_{\mathbf{MDP}} = \Sigma_{\mathcal{B}} + \Sigma_{\mathcal{W}_{\mathbb{R}}} + \Sigma_{\mathcal{R}_A} + \Sigma_{\diamond}, \quad \mathcal{U}_{\mathbf{MDP}} = ((\mathcal{B} \otimes \mathcal{U}_{\mathcal{W}_{\mathbb{R}}}) \otimes \mathcal{R}_A) + \mathcal{U}_{\diamond},$$

where $\mathcal{W}_{\mathbb{R}}$ is the theory of writing computations and the other theories are as in Section 8.

For convenience, we regard Markov decision processes over metric spaces as the coalgebras for the functor $(\Delta(\mathbb{R} \square c \cdot -))^A$ on \mathbf{Met} , where the endofunctor $(\mathbb{R} \square -)$ is used to encode the metric differences at each decision step for the real-valued reward available for two states. Via this coalgebraic representation, the c -discounted *probabilistic bisimilarity distance* on this structures can be defined as in [32] (following the same definition of Section 8).

► **Remark 34.** In [30] a Markov decision process is defined as a tuple $(S, p(\cdot|s, a), r(s, a))$ with a Markov kernel $p: S \times A \rightarrow \Delta(S)$ and randomised reward function $r: S \times A \rightarrow \Delta(\mathbb{R})$. Our coalgebraic representation is the natural generalisation over metric spaces, where the randomness of the Markov kernel and reward function is combined as a probability measure on $(\mathbb{R} \square c \cdot S)$, by regarding \mathbb{R} and S as extended metric spaces (for each $a \in A$).

On Metric Spaces and Complete Metric Spaces

Similarly to what we have done in Section 8 for labelled Markov processes, we relate Markov decision processes and their c -discounted probabilistic bisimilarity pseudometric with the free monads on the theory \mathcal{U}_{MDP} in **Met** and **CMet**.

The only step that changes in the characterisation of $T_{\mathcal{U}_{\text{MDP}}}$, regards the combination of theories $\mathcal{B} \otimes \mathcal{U}_{\mathcal{W}_{\mathbb{R}}}$, which is dealt using Corollary 28. Thus, similarly to Section 8 we get

$$T_{\mathcal{U}_{\text{MDP}}} = T_{((\mathcal{B} + \otimes \mathcal{U}_{\mathcal{W}_{\mathbb{R}}}) \otimes \mathcal{R}_A) + \mathcal{U}_{\circ}} \cong \mu y. \Pi((\mathbb{R} \square y) + -)^A.$$

The metric on the initial solution for the functorial fixed point definition corresponds to the c -discounted probabilistic bisimilarity (pseudo)metric on its coalgebra structure.

Similar considerations apply also when interpreting the theories in the category **CMet** of complete metric spaces, as the argument follows without issues because \mathbb{R} a complete metric space. Thus we obtain the following characterisation for the monad:

$$\mathbb{C}T_{\mathcal{U}_{\text{LMP}}} \cong \mu y. \Delta((\mathbb{R} \square y) + -)^A.$$

Again, the metric on the solution for the above functorial fixed point definition corresponds to the c -discounted probabilistic bisimilarity metric. Moreover, as the fixed point solution is unique, $\mathbb{C}T_{\mathcal{U}_{\text{LMP}}}0$ is an algebraic characterization of the final $(\Delta(\mathbb{R} \square c \cdot -))^A$ -coalgebra.

10 Conclusions

We studied the commutative combination of quantitative effects as the tensor of their quantitative equational theories. The key result in this regard is Theorem 17, asserting that the tensor of two quantitative theories corresponds to the categorical tensor of their free monads. In addition to this general result, we show how to extend to the quantitative algebraic setting Moggi's notions of reader and writer monad transformers.

We illustrate the applicability of our theoretical development with two examples: labeled Markov processes and Markov decision processes. Apart from the intrinsic interest in their quantitative equational axiomatisations, what is particularly pleasant is the systematic compositional way with which one can obtain quantitative axiomatisations of different variants of Markov processes by just combining theories as new basic ingredients.

An example that escapes our compositional treatment via sum and tensor is the combination of probabilities and non-determinism as illustrated in [23]. A possible future work in this direction is to extend the combination of theories with another operator: the distributive tensor (see [13, Section 6]). Following a similar intuition by Cheng [6], we claim that these correspond in a suitable way to Garner's weak distributive law [9]. Our claim seems promising in the light of the work [10, 3] which consider equational axiomatisations combining probabilities and non-determinism.

References

- 1 Giorgio Bacci, Radu Mardare, Prakash Panangaden, and Gordon D. Plotkin. An algebraic theory of markov processes. In *LICS*, pages 679–688. ACM, 2018.
- 2 Jon Beck. Distributive laws. In *Seminar on Triples and Categorical Homology Theory*, volume 80 of *Lect. Notes Math.*, pages 119–140. Springer, 1966.
- 3 Filippo Bonchi and Alessio Santamaria. Combining semilattices and semimodules. *CoRR*, abs/2012.14778, 2020.
- 4 Nathan J. Bowler, Sergey Goncharov, Paul Blain Levy, and Lutz Schröder. Exploring the boundaries of monad tensorability on set. *Log. Methods Comput. Sci.*, 9(3), 2013.
- 5 Pietro Cenciarelli and Eugenio Moggi. A syntactic approach to modularity in denotational semantics. Technical report, CWI, 1993. Proc. 5th. Biennial Meeting on Category Theory and Computer Science.
- 6 Eugenia Cheng. Distributive laws for lawvere theories. *Compositionality*, 2:1, May 2020. doi:10.32408/compositionality-2-1.
- 7 Josee Desharnais, Vineet Gupta, Radha Jagadeesan, and Prakash Panangaden. Metrics for labelled Markov processes. *Theoretical Computer Science*, 318(3):323–354, 2004.
- 8 Peter J. Freyd. Algebra valued functors in general and tensor products in particular. *Colloq. Math.*, 14:89–106, 1966.
- 9 Richard Garner. The vitoris monad and weak distributive laws. *Appl. Categorical Struct.*, 28(2):339–354, 2020.
- 10 Alexandre Goy and Daniela Petrisan. Combining probabilistic and non-deterministic choice via weak distributive laws. In *LICS*, pages 454–464. ACM, 2020.
- 11 Martin Hyland, Paul Blain Levy, Gordon D. Plotkin, and John Power. Combining algebraic effects with continuations. *Theor. Comput. Sci.*, 375(1-3):20–40, 2007.
- 12 Martin Hyland, Gordon D. Plotkin, and John Power. Combining effects: Sum and tensor. *Theor. Comput. Sci.*, 357(1-3):70–99, 2006. doi:10.1016/j.tcs.2006.03.013.
- 13 Martin Hyland and John Power. Discrete lawvere theories and computational effects. *Theor. Comput. Sci.*, 366(1-2):144–162, 2006.
- 14 Martin Hyland and John Power. The category theoretic understanding of universal algebra: Lawvere theories and monads. *Electronic Notes in Theor. Comp. Sci.*, 172:437–458, 2007.
- 15 Leonid Vitalevich Kantorovich. On the transfer of masses (in Russian). *Doklady Akademii Nauk*, 5(5-6):1–4, 1942. Translated in *Management Science*, 1958.
- 16 Gregory M. Kelly. Basic concepts of enriched category theory. *Theory and Applications of Categories*, 1982. Reprinted in 2005.
- 17 Anders Kock. Strong functors and monoidal monads. *Arch. Math. (Basel)*, 23:113–120, 1972.
- 18 Saunders Mac Lane. *Categories for the Working Mathematician*. Graduate Texts in Mathematics. Springer New York, 2nd edition, 1998.
- 19 William F. Lawvere. Metric spaces, generalized logic, and closed categories. In *Seminario Mat. e. Fis. di Milano*, volume 43, pages 135–166. Springer, 1973.
- 20 Ernest Manes. A Triple Theoretic Construction of Compact Algebras. In *Seminar on Triples and Categorical Homology Theory*, volume 80 of *Lect. Notes Math.*, pages 91–118. Springer, 1966.
- 21 Radu Mardare, Prakash Panangaden, and Gordon D. Plotkin. Quantitative Algebraic Reasoning. In *LICS*, pages 700–709. ACM, 2016. doi:10.1145/2933575.2934518.
- 22 Radu Mardare, Prakash Panangaden, and Gordon D. Plotkin. On the axiomatizability of quantitative algebras. In *LICS 2017*, pages 1–12. IEEE Computer Society, 2017. doi:10.1109/LICS.2017.8005102.
- 23 Matteo Mio and Valeria Vignudelli. Monads and quantitative equational theories for non-determinism and probability. In *CONCUR*, volume 171 of *LIPICs*, pages 28:1–28:18. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2020.
- 24 Eugenio Moggi. *The partial lambda calculus*. PhD thesis, University of Edinburgh. College of Science and Engineering. School of Informatics., 1988.

- 25 Eugenio Moggi. Notions of computation and monads. *Information and computation*, 93(1):55–92, 1991.
- 26 Gordon Plotkin and John Power. Semantics for algebraic operations. *Electronic Notes in Theoretical Computer Science*, 45:332–345, 2001.
- 27 Gordon Plotkin and John Power. Notions of computation determine monads. In *Foundations of Software Science and Computation Structures*, pages 342–356. Springer, 2002.
- 28 Gordon D. Plotkin and John Power. Semantics for algebraic operations. In *MFPS*, volume 45 of *Electronic Notes in Theoretical Computer Science*, pages 332–345. Elsevier, 2001.
- 29 Gordon D. Plotkin and John Power. Algebraic operations and generic effects. *Appl. Categorical Struct.*, 11(1):69–94, 2003.
- 30 M. L. Puterman. *Markov Decision Processes*. Wiley, 2005.
- 31 Marshall H. Stone. Postulates for the barycentric calculus. *Annali di Matematica Pura ed Applicata*, 29(1):25–30, 1949.
- 32 Franck van Breugel, Claudio Hermida, Michael Makkai, and James Worrell. Recursively defined metric spaces without contraction. *Theor. Comput. Sci.*, 380(1-2):143–163, 2007. doi:10.1016/j.tcs.2007.02.059.