

Positivity Problems for Reversible Linear Recurrence Sequences

George Kenison ✉

Institute of Logic and Computation, TU Wien, Austria

Joris Nieuwveld ✉

Max Planck Institute for Software Systems, Saarland Informatics Campus, Saarbrücken, Germany

Joël Ouaknine ✉

Max Planck Institute for Software Systems, Saarland Informatics Campus, Saarbrücken, Germany

James Worrell ✉

Department of Computer Science, University of Oxford, UK

Abstract

It is a longstanding open problem whether there is an algorithm to decide the Positivity Problem for linear recurrence sequences (LRS) over the integers, namely whether given such a sequence, all its terms are non-negative. Decidability is known for LRS of order 5 or less, i.e., for those sequences in which every new term depends linearly on the previous five (or fewer) terms. For *simple* LRS (i.e., those sequences whose characteristic polynomials have no repeated roots), decidability of Positivity is known up to order 9.

In this paper, we focus on the important subclass of *reversible* LRS, i.e., those integer LRS $\langle u_n \rangle_{n=0}^{\infty}$ whose bi-infinite completion $\langle u_n \rangle_{n=-\infty}^{\infty}$ also takes exclusively integer values; a typical example is the classical Fibonacci (bi-)sequence $\langle \dots, 5, -3, 2, -1, 1, 0, 1, 1, 2, 3, 5, \dots \rangle$. Our main results are that Positivity is decidable for reversible LRS of order 11 or less, and for simple reversible LRS of order 17 or less.

2012 ACM Subject Classification Mathematics of computing → Discrete mathematics; Computing methodologies → Algebraic algorithms

Keywords and phrases The Positivity Problem, Linear Recurrence Sequences, Verification

Digital Object Identifier 10.4230/LIPIcs.ICALP.2023.130

Category Track B: Automata, Logic, Semantics, and Theory of Programming

Funding *George Kenison*: Work supported by WWTF ICT19-018 grant ProbInG and ERC Consolidator Grant ARTIST 101002685.

Joël Ouaknine: Also affiliated with Keble College, Oxford as emmy.network Fellow, and supported by DFG grant 389792660 as part of TRR 248 (see <https://perspicuous-computing.science>).

James Worrell: Work supported by UKRI Frontier Research Grant EP/X033813/1.

1 Introduction

The Positivity Problem

The class of *threshold problems* considers whether a given loop program's variables remain above a fixed threshold before and after each iteration of the loop. In automated verification, this class of decision problems is relevant to program correctness, and particularly questions regarding termination, persistence, and reachability. The moniker *Positivity* is used when the chosen threshold is zero. In this paper, we shall consider the Positivity Problem (and its variants) for a particular class of integer-valued linear recurrence sequences.



© George Kenison, Joris Nieuwveld, Joël Ouaknine, and James Worrell;
licensed under Creative Commons License CC-BY 4.0

50th International Colloquium on Automata, Languages, and Programming (ICALP 2023).

Editors: Kousha Etessami, Uriel Feige, and Gabriele Puppis; Article No. 130; pp. 130:1–130:17

Leibniz International Proceedings in Informatics



LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany



130:2 Positivity Problems for Reversible Linear Recurrence Sequences

An integer-valued *linear recurrence sequence* (LRS) $\langle u_n \rangle_n$ satisfies a relation of the form

$$u_{n+d} = a_{d-1}u_{n+d-1} + \cdots + a_1u_{n+1} + a_0u_n \quad (1)$$

where the coefficients $a_{d-1}, \dots, a_1, a_0 \in \mathbb{Z}$ and, without loss of generality, we can assume that $a_0 \neq 0$. The sequence $\langle u_n \rangle_n$ is then wholly determined by the recurrence relation and the initial values u_0, u_1, \dots, u_{d-1} . The relation in (1) is said to have *length* d and the *order* of an LRS $\langle u_n \rangle_n$ is equal to the length of the shortest relation that $\langle u_n \rangle_n$ satisfies. The polynomial $f(X) = X^d - a_{d-1}X^{d-1} - \cdots - a_1X - a_0$ is the *characteristic polynomial* associated with relation (1).

Given an LRS $\langle u_n \rangle_n$, the *Positivity Problem* asks to determine whether $u_n \geq 0$ for each $n \in \mathbb{N}_0$. Positivity is a longstanding open problem and is intimately related to the well-known *Skolem Problem*, which asks to determine whether an LRS vanishes at some term [6, 8]. Indeed, if one could establish decidability of Positivity, then decidability of Skolem would necessarily follow (cf. [13]). One of the motivations to study Positivity lies in its connections to program verification [15]. Take, for example, the following linear loop P with inputs $\underline{w}, \underline{b} \in \mathbb{Z}^d$ and $A \in \mathbb{Z}^{d \times d}$ where

$$P: \underline{v} \leftarrow \underline{w}; \quad \mathbf{while} \quad \underline{b}^\top \cdot \underline{v} \geq 0 \quad \mathbf{do} \quad \underline{v} \leftarrow A\underline{v}. \quad (2)$$

Let $\langle u_n \rangle_n$ be the LRS with terms given by $u_n = \underline{b}^\top A^n \underline{w}$. It is clear that loop P terminates if and only if there exists an $n \in \mathbb{N}_0$ such that $u_n < 0$. Conversely, to each LRS $\langle u_n \rangle_n$ we can associate a linear loop of the form (2): one need only take A to be the transpose of the companion matrix associated with $\langle u_n \rangle_n$ so that

$$A = \begin{pmatrix} a_{d-1} & 1 & \cdots & 0 & 0 \\ \vdots & 0 & \ddots & 0 & 0 \\ a_2 & 0 & \cdots & 1 & 0 \\ a_1 & 0 & \cdots & 0 & 1 \\ a_0 & 0 & \cdots & 0 & 0 \end{pmatrix}, \quad \underline{b}^\top = (u_{d-1}, \dots, u_1, u_0), \quad \text{and} \quad \underline{w} = (1, 0, \dots, 0)^\top$$

in order to recover the terms $u_{n+d-1} = \underline{b}^\top A^n \underline{w}$.

Variants of the Positivity Problem have also garnered attention in the literature. For example, the *Ultimate Positivity Problem* weakens the guard clause: given an LRS $\langle u_n \rangle_n$, determine whether there exists an $N \in \mathbb{N}$ such that $u_n \geq 0$ if $n > N$. By contrast, the *Simple Positivity Problem* restricts the class of sequences under consideration to those that are simple. Here an LRS $\langle u_n \rangle_n$ is *simple* if each of the roots of the associated characteristic polynomial has algebraic multiplicity one. In this paper, we focus on the *Reversible Positivity Problem*, i.e., the restriction of the Positivity Problem to the class of LRS that are reversible (as defined below).

Background and Motivation

Lipton *et al.* [11] coined the term *reversible* to describe the class of LRS that assume exclusively integer values, whether the sequences are expanded forwards or backwards. Equivalently, such LRS can be shown to satisfy relations of the form (1) with the condition $a_0 = \pm 1$ (or, alternatively, the associated characteristic polynomial satisfies $f(0) = \pm 1$). The subclass of while loops (as in (2)) naturally associated with reversible sequences have *unimodular* update matrices. The inverse A^{-1} of a unimodular matrix A is likewise unimodular. Thus the uniquely defined bi-infinite extension $\langle u_n \rangle_{n=-\infty}^\infty$ with each $u_n = \underline{b}^\top A^n \underline{w}$ (as above) is integer-valued.

The decidability of Reversible Skolem – where one restricts the Skolem Problem to reversible LRS – was established up to order 7 in a recent paper by Lipton *et al.* [11]. Kenison [9] gave an alternative proof of this decidability result, leveraging a powerful result for algebraic units due to Dubickas and Smyth [5]. In general, the Skolem Problem is only known to be decidable for arbitrary LRS up to order four [12, 22]. The Positivity Problem is decidable for arbitrary LRS of order five or less [16], and for simple LRS of order 9 or less [13]. The Ultimate Positivity Problem is also decidable for arbitrary LRS of order 5 or less, as well as for simple LRS of arbitrary order [14, 16]. Previous work showed that the Positivity Problem is decidable for simple reversible LRS of order 10 or less [9].

Contributions

In this paper, we shall consider Positivity problems for reversible LRS. We will exploit spectral properties of reversible LRS and employ techniques from both Galois theory and Diophantine approximation to establish decidability at higher orders than is currently known for general positivity. Our main contributions are as follows:

- ▶ **Theorem 1.** *For reversible LRS, the Positivity and Ultimate Positivity Problems are both decidable up to order 11.*
- ▶ **Theorem 2.** *For simple reversible LRS, the Positivity Problem is decidable up to order 17.*

Structure

The remainder of this paper is structured as follows. In the next section we review necessary preliminary material. In Section 3, we prove results on the root structures of characteristic polynomials associated with reversible LRS. In Section 4, we prove Theorems 1 and 2. We also consider barriers impeding further progress to the state of the art (i.e., decidability results at higher orders) by exhibiting sequences that are not amenable to standard Diophantine approximation techniques due to certain spectral properties (see Section 5). The calculations involved in preparing these hard instances were performed in SageMath [4].

2 Preliminaries

2.1 Linear Recurrence Sequences

We expand upon the standard terminology for LRS given in the introduction. It is straightforward to see that an LRS $\langle u_n \rangle_n$ is wholly determined by a recurrence relation (as in (1)) and the initial values u_0, u_1, \dots, u_{d-1} .

Let $\langle u_n \rangle_n$ be an LRS with characteristic polynomial f . It is well-known that an LRS admits a closed-form representation as an exponential polynomial; specifically, for each $n \in \mathbb{N}_0$, we have $u_n = A_1(n)\lambda_1^n + \dots + A_\ell(n)\lambda_\ell^n$. Here the *characteristic roots* $\lambda_1, \dots, \lambda_\ell$ are the distinct roots of f . Further, the polynomial coefficients $A_j \in \overline{\mathbb{Q}}[X]$ are completely determined by the initial values of $\langle u_n \rangle_n$. The polynomial coefficients for a simple LRS are all constants; that is, if $\langle u_n \rangle_n$ is simple, then $u_n = A_1\lambda_1^n + \dots + A_\ell\lambda_\ell^n$.

An LRS is *degenerate* when there are two distinct characteristic roots whose quotient is a root of unity. Otherwise, the sequence is said to be *non-degenerate*.

Let $\lambda_1, \dots, \lambda_\ell$ be the characteristic roots of an LRS $\langle u_n \rangle_n$. A characteristic root λ of $\langle u_n \rangle_n$ is *dominant* if $|\lambda| \geq |\lambda_j|$ for each $j \in \{1, \dots, \ell\}$. By convention, when we talk about the number of dominant roots *we do not count multiplicity*; e.g., a recurrence sequence that satisfies the relation $u_{n+2} = 2u_{n+1} - u_n$ with characteristic polynomial $(X - 1)^2$ has only one dominant root.

In the sequel, we will state and prove technical results for polynomials in $\mathbb{Z}[X]$. Here we say that a polynomial in $\mathbb{Z}[X]$ is *non-degenerate* if no quotient of distinct roots is a root of unity and we say that it is *reversible* if it is monic and has constant term ± 1 . Note that our use of these terms for both recurrence sequences and their characteristic polynomials is consistent.

2.2 The Positivity Problem

In this subsection we briefly recall decidability results for the Positivity Problem for LRS. We first recall the standard assumptions that permit us to reduce the problem of deciding positivity to that of deciding positivity for LRS that are both non-degenerate and possess a positive dominant characteristic root.

First, it is well known (cf. [6, 8]) that we can effectively reduce the computational study of LRS to that of non-degenerate LRS. This observation follows because each degenerate LRS can be realised as an interleaving of finitely many non-degenerate LRS of the same order. Thus we need only consider non-degenerate LRS when studying positivity. We note also that this reduction preserves the quality of having simple characteristic roots.

Second, let us recall the following classical consequence of the Vivanti–Pringsheim Theorem from complex analysis [17, 23] (see also [21, Section 7.21]).

► **Lemma 3.** *Suppose that a non-zero LRS $\langle u_n \rangle_n$ has no positive dominant characteristic root. Then the sets $\{n \in \mathbb{N} : u_n > 0\}$ and $\{n \in \mathbb{N} : u_n < 0\}$ are both infinite.*

As a consequence of Lemma 3, we can reduce the problem of deciding positivity to LRS that possess a positive dominant characteristic root.

Together, the two preceding assumptions show that the sequences we consider have an odd number of dominant roots: the set of dominant roots comprises complex-conjugate pairs of roots and a single positive dominant root. Note that a second real dominant root would violate non-degeneracy.

Ouaknine and Worrell showed that the Simple Positivity Problem for LRS is decidable up to order nine [13]. The main technical contribution of that paper was the following result, which, in combination with the various observations above, covers all sequences up to order nine.

► **Theorem 4.** *Let $\langle u_n \rangle_n$ be a non-degenerate simple LRS with characteristic polynomial $f \in \mathbb{Z}[X]$ and a positive dominant root. If $f \in \mathbb{Z}[X]$ has either at most eight dominant roots or precisely nine roots, then we can determine whether $u_n \geq 0$ for each $n \in \mathbb{N}_0$.*

2.3 Number Theory

An algebraic integer is a *unit* if its multiplicative inverse is also an algebraic integer. It is a basic fact that an algebraic number is a unit if and only if its minimum polynomial in $\mathbb{Z}[X]$ is monic and has constant term ± 1 . Thus the characteristic roots of a reversible LRS are all units.

A *number field* K is a field determined by a finite extension of \mathbb{Q} . The *splitting field* K for the polynomial $f \in \mathbb{Q}[X]$ is the field extension of \mathbb{Q} with the following two properties. First, the polynomial f can be written as a product of linear factors in $K[X]$ (i.e., f splits completely in K) and second, f does not split completely over any proper subfield of K containing \mathbb{Q} .

2.4 Group Theory

A finite group G is said to act *transitively* on a finite set X if for each pair $x, y \in X$ there is a $g \in G$ such that $gx = y$. The *stabilizer* G_x of an element x in X is defined as the set $\{g \in G : gx = x\}$. The Orbit-Stabilizer Theorem (see, for example, [18, Theorem 3.19]) implies that if G acts transitively on X , the cardinality of G_x is the same for each $x \in X$. Further, $\#\{g \in G : gx = y\}$ is the same for all $x, y \in X$ in this scenario.

2.5 Galois Theory

We assume familiarity with basic notions in Galois theory and the theory of number fields. For reference, we recommend standard textbooks such as [3, 20]. The following includes some of the definitions and theory we use in the sequel.

Given a field extension K of \mathbb{Q} , we use $\text{Gal}_{\mathbb{Q}}(K)$ to denote the *Galois group of K over \mathbb{Q}* ; that is, the group of automorphisms of K that fix \mathbb{Q} . We shall refer to elements of a Galois group as *Galois automorphisms*. In the sequel, we shall frequently use the following property of irreducible polynomials. Let $f \in \mathbb{Q}[X]$ be an irreducible polynomial and K the splitting field of f . Then the Galois group $\text{Gal}_{\mathbb{Q}}(K)$ acts transitively on the roots of f . Indeed, the *orbit* of a root of f (i.e., the set of images of the root under the group action) is the set of roots of f . In light of the above, for a given algebraic number α we use the term *Galois conjugates* to refer to set of roots of the minimal polynomial of α .

Recall the following theorem due to Kronecker [10].

► **Theorem 5.** *Let $f \in \mathbb{Z}[X]$ be a monic polynomial such that $f(0) \neq 0$. Suppose that all the roots of f have absolute value at most 1. Then all the roots of f are roots of unity.*

We deduce the following. If $f \in \mathbb{Z}[X]$ is the characteristic polynomial of a reversible LRS $\langle u_n \rangle_n$ such that the roots of f all lie in the unit disk $\{z \in \mathbb{C} : |z| \leq 1\}$, then the roots of f are all roots of unity and so $\langle u_n \rangle_n$ is of order one or degenerate. In the former case, positivity of $\langle u_n \rangle_n$ is easily determined and in the latter case, determining whether $\langle u_n \rangle_n$ is positive reduces to studying positivity for associated non-degenerate LRS. Thus in the sequel we shall always assume, without loss of generality, that the dominant roots of f lie on a circle with radius strictly larger than 1.

The roots of an irreducible polynomial are necessarily Galois conjugates. We call the quotient of two distinct roots of an irreducible polynomial a *conjugate ratio*.

Key to the technical lemmas we prove in the sequel are results concerning identities between the roots of irreducible polynomials. We employ a powerful result due to Dubickas and Smyth [5], Theorem 6 below, concerning necessary conditions for an algebraic unit and all its Galois conjugates to lie on two concentric circles centred at the origin. (Theorem 6 is a specialisation of the general result [5, Theorem 2.1].)

► **Theorem 6.** *Suppose that $f \in \mathbb{Z}[X]$ is an irreducible, reversible polynomial of degree d such that all the roots of f lie on two circles centred at the origin. Let r and R be the radii of the respective circles and, without loss of generality, suppose that at most half of the roots of f lie on the circle of radius r . Then we have the following: either d is even, in which case half of the roots lie on the circle of radius r ; or d is a multiple of three and a third of the roots lie on the circle of radius r . In the latter case, we additionally have that for every root β on the circle of radius r there exists $n > 0$ such that $\beta^n \in \mathbb{R}$.*

We shall frequently employ the following lemma, versions of which were proved by Smyth [19] and Ferguson [7].

► **Lemma 7.** *Suppose that λ is an algebraic number with Galois conjugates β and γ satisfying $\lambda^2 = \beta\gamma$. Then the conjugate ratio λ/β is a root of unity.*

3 Root Analysis of Reversible Polynomials

The main result of this section is Theorem 14, concerning the number of dominant roots of a reversible polynomial. Essentially the theorem says that, excepting a number of special cases, no more than half of the roots of such a polynomial can be dominant. This is the key technical tool behind our main decidability results for the Positivity Problem for reversible LRS.

Let us begin with two lemmas concerning the dominant roots of reversible polynomials. These can be considered as weak forms of the main result of the section (and are used in the proof thereof).

► **Lemma 8.** *Let $f \in \mathbb{Z}[X]$ be an irreducible non-degenerate polynomial with a real dominant root λ . Then f has exactly one dominant root.*

Proof. Let λ be a real dominant root. Suppose β is also a dominant root. Then $\lambda^2 = \beta\bar{\beta}$ and hence λ/β is a root of unity by Lemma 7. Since f is non-degenerate we have $\lambda = \beta$; that is, f has exactly one dominant root. ◀

► **Lemma 9.** *Suppose that $f \in \mathbb{Z}[X]$ is irreducible, non-degenerate, and reversible, with $2m$ non-real dominant roots and no real dominant roots. Then $\deg(f) > 3m$ if $m \geq 2$. Further, $\deg(f) \leq 3m$ only if $(\deg(f), m) = (3, 1)$ or f is constant.*

Proof. Since f has at least $2m$ roots, it is clear that $\deg(f) \geq 2m$. The case where $m = 0$ pertains to constant polynomials, thus we need only consider the case when $m \geq 1$.

We will first show that $\deg(f) > 2m$. Assume, for a contradiction, that $\deg(f) = 2m$. Then the roots of f all lie on the circumference of some circle centred at the origin. We make the following two observations. First, f is reversible, and hence monic. Second, by Vieta's formulas, $|f(0)| = 1$ is equal to the absolute value of the product of the roots of f . From these observations, we conclude that the roots of f all lie on the unit circle and, by Theorem 5, are therefore roots of unity. As $m \geq 1$, f has at least two roots, and their conjugate ratio is thus a root of unity. We have reached a contradiction: f is assumed to be non-degenerate. Thus $\deg(f) > 2m$.

Consider the subcase where $m = 1$. Assume that $2m < \deg(f) \leq 3m$, then clearly we have $\deg(f) = 3$. The assertion in the lemma trivially holds. Hereafter we assume that $m \geq 2$.

We now show that under the assumption that $m \geq 2$, we necessarily have $\deg(f) \geq 3m$. Let $\lambda_1, \bar{\lambda}_1, \dots, \lambda_m, \bar{\lambda}_m$ be the $2m$ dominant roots of f . Thus $\lambda_1 \bar{\lambda}_1 = \lambda_i \bar{\lambda}_i$ for each $i \in \{1, \dots, m\}$. Since $2m < \deg(f)$, f has a non-dominant root γ . Further, since f is irreducible, there is a Galois automorphism σ such that $\sigma(\lambda_1) = \gamma$. We claim that for each $i \in \{2, \dots, m\}$ at least one of $\sigma(\lambda_i)$ and $\sigma(\bar{\lambda}_i)$ is a non-dominant root of f . Assume, for a contradiction, that the claim does not hold. Then there is an $i \in \{2, \dots, m\}$ such that both $\sigma(\lambda_i)$ and $\sigma(\bar{\lambda}_i)$ are dominant roots. The map σ necessarily preserves polynomial symmetries between the roots of f and so $\gamma\sigma(\bar{\lambda}_1) = \sigma(\lambda_i)\sigma(\bar{\lambda}_i)$. However, since $\sigma(\lambda_1) = \gamma$ is strictly smaller in absolute value than both $\sigma(\lambda_i)$ and $\sigma(\bar{\lambda}_i)$, we have $|\gamma\sigma(\bar{\lambda}_1)| < |\sigma(\lambda_i)\sigma(\bar{\lambda}_i)|$. This last inequality contradicts the aforementioned symmetry between dominant roots. We conclude that the list of non-dominant roots of f includes γ and at least one of $\sigma(\lambda_i)$ and $\sigma(\bar{\lambda}_i)$ for each $i \in \{2, \dots, m\}$. Thus f has at least m non-dominant roots and so $\deg(f) \geq m + 2m = 3m$.

Finally, we eliminate the case that $\deg(f) = 3m$ when $m \geq 2$. Assume, for a contradiction, that $\deg(f) = 3m$. We can apply the preceding argument to the reciprocal polynomial of f to deduce that the m non-dominant roots of f are equal in modulus and so all lie on a circle $\{z \in \mathbb{C} : |z| = r\}$ for some $r > 0$. Thus all roots of the irreducible and reversible polynomial

130:8 Positivity Problems for Reversible Linear Recurrence Sequences

Proof. Assume, for a contradiction, that the conjugate ratio $\mu_j/\mu_{j'}$ of g is a root of unity. Both root sets \mathcal{A}_j and $\mathcal{A}_{j'}$ have cardinality $2m$. Since $\deg(f) < 4m = \#\mathcal{A}_j + \#\mathcal{A}_{j'}$, we deduce that $\mathcal{A}_j \cap \mathcal{A}_{j'}$ is non-empty. Let $\lambda \in \mathcal{A}_j \cap \mathcal{A}_{j'}$ and κ, κ' be roots of f such that $\lambda\kappa = \mu_j$ and $\lambda\kappa' = \mu_{j'}$. Since $\mu_j \neq \mu_{j'}$, we have $\kappa \neq \kappa'$. It follows that f is degenerate because $\kappa/\kappa' = \mu_j/\mu_{j'}$ is a root of unity. From this contradiction, we deduce that g is non-degenerate. \blacktriangleleft

► **Lemma 11.** *Suppose that $f \in \mathbb{Z}[X]$ is reversible, non-degenerate, and irreducible with $2m$ non-real dominant roots. Write g for the dominant polynomial of f . Then all the roots of f have the same equation number E and*

$$2m \deg(g) = E \deg(f). \quad (4)$$

Proof. We use the notation of $\lambda_i, \mu_j, \sigma_j, \tilde{\sigma}_j, \alpha_{i,j,k}, K, L$, etc. as above.

Set $H = \text{Gal}_{\mathbb{Q}}(K)$ and $G = \text{Gal}_{\mathbb{Q}}(L)$. By the Orbit-Stabilizer Theorem, the number of $\sigma \in H$ such that $\sigma(\mu_1) = \mu_j$ is independent of the choice of $j \in \{1, \dots, n\}$. Now each $\sigma \in H$ has the same number of lifts to G , and so the number of elements of G that map μ_1 to each μ_j is independent of $j \in \{1, \dots, n\}$. Thus the number of elements of G such that the image of \mathcal{A}_1 is \mathcal{A}_j is also independent of the choice of j .

We make the following claim whose proof is given immediately below.

▷ **Claim 12.** In the setting defined above, there is no pair of distinct j_1 and j_2 for which $\mathcal{A}_{j_1} = \mathcal{A}_{j_2}$.

We also make the following observation. By the Orbit-Stabilizer Theorem, for every choice of two roots λ and λ' of f , the number of $\sigma \in G$ such that $\sigma(\lambda) = \sigma(\lambda')$ is equal. Thus for each root λ of f , the number of $\sigma \in G$ such that one of $\tilde{\sigma}(\lambda_1), \tilde{\sigma}(\overline{\lambda_1}), \dots, \tilde{\sigma}(\lambda_m), \tilde{\sigma}(\overline{\lambda_m})$ equals λ is independent of the choice of λ . This shows that the equation number E is independent of the choice of the root λ .

The equation $2m \deg(g) = E \deg(f)$ follows from counting the number of $\alpha_{i,j,k}$. On the one hand, there are $\deg(g)$ equations with $2m$ entries (Claim 12). On the other hand, there are $\deg(f)$ roots each appearing E times. \blacktriangleleft

Proof of Claim 12. Let us assume, for a contradiction, that $\mathcal{A}_{j_1} = \mathcal{A}_{j_2}$ for $j_1 \neq j_2$. Then $\mu_{j_1} \neq \mu_{j_2}$ and

$$\mu_{j_1}^m = \prod_{i=1}^m \alpha_{i,j_1,1} \alpha_{i,j_1,2} = \prod_{i=1}^m \alpha_{i,j_2,1} \alpha_{i,j_2,2} = \mu_{j_2}^m.$$

Thus μ_{j_1}/μ_{j_2} is a root of unity. Since $\alpha_{1,j_1,1} \in \mathcal{A}_{j_2}$, there are $1 \leq i \leq m$ and $k \in \{1, 2\}$ such that $\alpha_{1,j_1,1} = \alpha_{i,j_2,k}$. Then we have that the conjugate ratio $\alpha_{1,j_1,1}/\alpha_{i,j_2,3-k}$ given by

$$1 \neq \frac{\mu_{j_1}}{\mu_{j_2}} = \frac{\alpha_{1,j_1,1} \alpha_{1,j_1,2}}{\alpha_{i,j_2,1} \alpha_{i,j_2,2}} = \frac{\alpha_{1,j_1,2}}{\alpha_{i,j_2,3-k}}$$

is also a root of unity. Since $\alpha_{1,j_1,2}$ and $\alpha_{i,j_2,3-k}$ are distinct roots of f whose quotient is a root of unity, it follows that f is degenerate. We have reached a contradiction to our assumption that f is non-degenerate. Thus we have the claimed result. \blacktriangleleft

The next result increases the bound on the degree of f to $\deg(f) \geq 4m$.

► **Theorem 13.** *Let $f \in \mathbb{Z}[X]$ be an irreducible, non-degenerate, and reversible polynomial with $2m$ dominant non-real roots and no real dominant roots, then $(\deg(f), m) = (3, 1)$ or $\deg(f) \geq 4m$.*

Proof. Assume, for a contradiction, that f is a minimal counterexample in the sense that all polynomials of strictly smaller degree satisfy the statement in Theorem 13.

From Lemma 9, we deduce that $\deg(f) > 3m$ if we are not in the exceptional case $(\deg(f), m) = (3, 1)$. As we assume that f is a counterexample to Theorem 13, $\deg(f) < 4m$ as well. We shall employ the preceding notation for the dominating polynomial g , the sets of roots \mathcal{A}_j of f , and the equation number E .

Consider that there are $2m$ distinct roots of f in each equation in (3). Since $\deg(f) < 4m$ and f has $2m$ dominant roots, there is at least one dominant root of f in each such equation. Let γ be a root of f with minimal absolute value, then $|\gamma\lambda_1|$ is the minimal absolute value attained by any root of g . We now show that at least half of the roots of g lie on the circle $\{z \in \mathbb{C} : |z| = |\gamma\lambda_1|\}$. Observe that γ is witnessed in E (and so more than half) of the pairings in (3) and, further, is necessarily paired with a dominant root (for otherwise, a pairing between γ and a non-dominant root breaks the equality in (3)). From (4) and our assumption that $\deg(f) < 4m$, we deduce that $2E > \deg(g)$, and so γ appears in more than half of the equations in (3). Each such equation is in correspondence with a root of g of minimal absolute value.

Consider the polynomial $h(X) := g(0)X^n g(X^{-1})$. The polynomial $X^n g(X^{-1})$ is the reciprocal polynomial of g and so immediately, the roots of h are precisely $\mu_1^{-1}, \dots, \mu_n^{-1}$ and $n = \deg(g) = \deg(h)$. From the preceding discussion, more than half of the roots of h are dominant. Moreover, we can easily deduce that h is reversible, irreducible, and non-degenerate as g has these properties. Thus h is another counterexample to the statement in Theorem 13. All that remains is to derive a contradiction from our assumption that f has minimal degree. We derive this contradiction by proving that $\deg(h) < \deg(f)$ and h does not belong to either one of the exceptional cases.

Observe that h cannot belong to one of the exceptional cases since (4) has no integer solutions when $n = 1, 2, 3$ and $3m < \deg(f) < 4m$. Thus it remains to show that $n \geq \deg(f)$ is absurd. Let us assume, for a contradiction, that λ_1 appears in a product pair with a dominant root other than $\bar{\lambda}_1$ in the j th equation of (3). Then μ_1 and μ_j have equal absolute value. If μ_j is real, $\mu_1/\mu_j = \pm 1$ contradicting our non-degeneracy assumption (Lemma 10). Similarly, we derive a contradiction to our non-degeneracy assumption if μ_j is non-real by Lemma 7. Thus, we can pair λ_1 with the $\deg(f) - 2m < 2m$ non-dominant roots of f and $\bar{\lambda}_1$. This gives the upper bound $E \leq \deg(f) - 2m + 1 \leq 2m$ on E . We substitute our assumption that $n = \deg(g) \geq \deg(f)$ into (4) to obtain a lower bound $2m \leq E$. Thus, $E = 2m$.

We use the equality $E = 2m$ to deduce that $\deg(f) = 4m - 1$ and make the following observations. Each of the $2m$ dominant roots of f pair with their respective complex conjugate and all of the $2m - 1$ non-dominant roots of f . Thus we can pair each non-dominant root of f with $2m$ dominant roots. Further, every pair of non-dominant roots of f appears in at least one equation in (3). Thus the roots of g and h lie on two concentric circles centred at the origin. The roots of g are distributed so that g has exactly one dominant root and $2m + (2m - 1) - 1 = 4m - 2$ non-dominant roots. By construction, h has exactly one non-dominant root and $4m - 2$ dominant roots. This distribution of roots is not possible by Theorem 6, hence a contradiction.

In summary, f cannot be a counterexample to Theorem 13 of smallest possible degree. We thus deduce that all polynomials that satisfy the hypothesis in the theorem obey the bound $\deg(f) \geq 4m$, as required. \blacktriangleleft

The only superfluous assumption in Theorem 13 is that f is irreducible. We circumvent the irreducibility assumption with a careful case analysis.

► **Theorem 14.** *Let f be a non-degenerate reversible polynomial. Suppose that more than half of the roots of f are dominant. Then either f is linear or f is cubic with two dominant roots.*

Proof. Let f be a counterexample of minimal degree, and factor f into irreducible polynomials f_1, \dots, f_k . For $1 \leq i \leq k$, let m'_i be the number of dominant roots of f_i . Call an irreducible factor *sharp* if $2m'_i = \deg(f_i)$ and *special* if $2m'_i > \deg(f_i)$. From Lemma 8 and Theorem 13, it follows that if an irreducible factor is special, then $(\deg(f_i), m'_i) = (1, 1)$ or $(3, 2)$. If $k = 1$, then f is irreducible and the result follows automatically. Thus we can freely assume that $k \geq 2$. Since f is a counterexample of minimal degree, a straightforward proof by contradiction permits us to assume $k = 2$. Thus our argument reduces to the following cases: we need only show that the product of either two special polynomials or a special and a sharp polynomial breaks the hypothesis. By renumbering, we can assume f_1 is special and f_2 is either sharp or special. We observe that under our assumptions the dominant roots of f_1 and f_2 are necessarily equal in absolute value and, as we do not count multiplicity, $f_1 \neq f_2$.

We begin our case analysis. First, consider the case where $(\deg(f_1), m'_1) = (1, 1)$. Then $f_1(X) = X \pm 1$ as f_1 is reversible. Since the root ∓ 1 of f_1 is a dominant root of f , we deduce that all roots of f lie on the unit circle as the roots of f are algebraic units. When we combine Lemma 8, Theorem 13, and our assumption that at least half of the roots of f are dominant, we deduce that $(\deg(f_2), m'_2) = (1, 1)$ and so $f_2(X) = X \mp 1$. Thus -1 and 1 are both roots of f , which contradicts our assumption that f is non-degenerate.

Second, let us suppose that $(\deg(f_1), m'_1) = (3, 2)$. Following the argument in the preceding case, either $(\deg(f_2), m'_2) = (3, 2)$ or $\deg(f_2) = 2m'_2$. In the former, the non-dominant roots γ_1 and γ_2 of f_1 and f_2 (respectively) are both real and equal in modulus. This is straightforward to see since each f_j is of the form $f_j = (x - \gamma_j)(x - Re^{i\theta_j})(x - Re^{-i\theta_j})$ for some $R > 1$ and $\gamma_j := \pm R^{-2}$. We cannot have two such irreducible factors since then the ratio $\gamma_1/\gamma_2 = \pm 1$, which breaks either the non-degeneracy assumption on f or the assumption that $f_1 \neq f_2$.

We continue with the latter subcase $(\deg(f_1), m'_1) = (3, 2)$ and $\deg(f_2) = 2m'_2$. Since the dominant roots of f_1 and f_2 are dominant roots of f , the dominating polynomials of f_1 and f_2 are one and the same, say g . Let E_1 and E_2 be the respective equation numbers of f_1 and f_2 . From (4), $2 \deg(g) = E_1 \deg(f_1) = 3E_1$. We thus deduce that E_1 is even. Since $1 \leq E_1 \leq \deg(f_1) = 3$ (each pairing is distinct), we have that $E_1 = 2$ and, it follows immediately, $\deg(g) = 3$. We substitute this result and our assumption that $\deg(f_2) = 2m'_2$ into (4) in order to obtain $m'_2 \deg(g) = 3m'_2 = 2E_2 m'_2$. We have reached a contradiction: $E_2 = 3/2$ is not an integer. We have exhausted the possibilities for constructing a minimal counterexample f and find that no such counterexample exists. We have thus proved Theorem 14. ◀

4 Decidability at Low Orders

In this section we complete the proofs of our two main theorems concerning the Positivity Problem for reversible LRS. We start with Theorem 2, which states that positivity of reversible sequences that are moreover simple is decidable up to order 17.

Proof of Theorem 2. As previously noted, we can reduce the Simple Reversible Positivity Problem to deciding positivity for the subclass of simple reversible LRS that are additionally both non-degenerate and in possession of a positive dominant root.

In light of the preceding paragraph, consider the subclass of non-degenerate, simple, and reversible LRS with a positive dominant root. Let f be the characteristic polynomial associated with a sequence in this class. Without loss of generality, we can additionally

assume that fewer than half of the roots of f are dominant. If $f \in \mathbb{Z}[X]$ has at least nine dominant roots, then, by Theorem 14, we have the bound $\deg(f) \geq 18$. Taking the contrapositive, if f is again the characteristic polynomial of a sequence in this subclass with $\deg(f) \leq 17$, then f has at most eight dominant roots.

Now we invoke Theorem 4 to deduce that, in the aforementioned subclass, positivity is decidable for LRS up to order 17. As noted at the beginning of this proof, this deduction is sufficient to obtain the desired result: simple reversible positivity is decidable up to order 17. ◀

We now turn our attention to general reversible sequences; i.e., we no longer assume that the characteristic roots are simple. Here, as stated in Theorem 1, we have decidability up to order 11.

Proof of Theorem 1. Assume, for a contradiction, that $\langle u_n \rangle_n$ is a reversible LRS and counterexample to the statement; that is to say, $\langle u_n \rangle_n$ is a reversible LRS with order at most 11 for which we cannot determine positivity or ultimate positivity.

From our earlier discussion on the Positivity Problem and Ultimate Positivity Problem in Subsection 2.2, it follows that we can reduce both problems for reversible LRS to deciding (ultimate) positivity for the subclass of reversible LRS that are additionally both non-degenerate and in possession of a positive dominant root.

For the class of reversible LRS with one dominant root, decidability of (ultimate) positivity is considered folklore. Thus we freely assume that $\langle u_n \rangle_n$ has at least three dominant roots (the positive root and a pair of complex conjugate roots). By Theorem 2 for positivity and the earlier mentioned results in [14] for ultimate positivity, we can also assume that $\langle u_n \rangle_n$ has a non-simple characteristic root. Now consider the exponential polynomial representation of $\langle u_n \rangle_n$: deciding (ultimate) positivity for LRS whose dominant characteristic roots are all simple reduces to deciding (ultimate) positivity for simple LRS. So, in addition, we shall assume that $\langle u_n \rangle_n$ has a non-simple dominant characteristic root. We will use the following claims, whose proofs are given immediately below.

▷ **Claim 15.** Suppose that the real positive dominant root ρ of sequence $\langle u_n \rangle_n$ (as above) is the only non-simple dominant root of $\langle u_n \rangle_n$. Then we can determine whether $\langle u_n \rangle_n$ is (ultimately) positive.

▷ **Claim 16.** Suppose that sequence $\langle u_n \rangle_n$ (as above) possesses non-real dominant roots that are not simple and, further, that the real dominant root ρ is simple. Then $\langle u_n \rangle_n$ is neither ultimately positive nor positive.

In light of the preceding claims, we freely assume that the counterexample $\langle u_n \rangle_n$ has at least three non-simple dominant characteristic roots and this collection must include the real dominant root ρ as well as a complex conjugate pair λ and $\bar{\lambda}$.

Let f be the monic integer-valued polynomial of the smallest degree with ρ and λ as roots. Then, f is non-degenerate and reversible. By Theorem 14, it follows that at most half of the roots of f are dominant if f is neither linear nor cubic with two dominant roots. As such, f has degree at least 6 and, additionally, as each of these roots is non-simple being a Galois conjugate of either ρ or λ , $\langle u_n \rangle_n$ has order at least 12.

We thus deduce the desired result: positivity and ultimate positivity are decidable for sequences up to order 11. ◀

130:12 Positivity Problems for Reversible Linear Recurrence Sequences

Proof of Claim 15. Suppose that the real positive dominant root ρ of $\langle u_n \rangle_n$ is the only non-simple dominant root of the sequence. If such a phenomenon were to take place then $u_n = A_\rho(n)\rho^n + O(\rho^n)$ where A_ρ is a non-constant polynomial. It is straightforward to deduce that $\langle u_n \rangle_n$ is (ultimately) positive if and only if $A_\rho(n)$ is (ultimately) positive in this instance. \triangleleft

Proof of Claim 16. We will show that the claim follows from Lemma 17 (cf. [1]).

► **Lemma 17.** *Let $\gamma_1, \dots, \gamma_k \in \{z \in \mathbb{C} : |z| = 1, z \neq 1\}$ be distinct complex numbers, $\alpha_1, \dots, \alpha_k \in \mathbb{C} \setminus \{0\}$, and $w_n = \sum_{\ell=1}^k \alpha_\ell \gamma_\ell^n$. Then there is a $c < 0$ such that $\operatorname{Re}(w_n) < c$ for infinitely many n .*

To prove the claim, let d be the maximum of the degrees of the roots of $\langle u_n \rangle_n$. Note $d \geq 1$ since, by assumption, $\langle u_n \rangle_n$ has a non-real dominant root that is not simple. We consider the normalised sequence $\langle v_n \rangle_n$ with terms given by $v_n = u_n/(\rho^n n^d)$ where ρ is the dominant root of $\langle u_n \rangle_n$. Note it is sufficient to establish that $\langle v_n \rangle_n$ is neither positive nor ultimately positive to obtain the desired result.

An analysis of the exponential polynomial of $\langle u_n \rangle_n$ leads to

$$v_n = \sum_{\ell=1}^{2k} \frac{A_\ell(n)}{n^d} \frac{\lambda_\ell^n}{\rho^n} + O(n^{-d})$$

where $\lambda_1, \dots, \lambda_{2k}$ are the non-real dominant roots of $\langle u_n \rangle_n$ and the implied constant associated with $O(n^{-d})$ is real-valued. Let α_ℓ be the leading coefficient of $A_\ell(n)$. Then $A_\ell(n)/n^d \rightarrow \alpha_\ell$ as $n \rightarrow \infty$. Now

$$v_n < r(n) + \sum_{\ell=1}^{2k} \alpha_\ell \frac{\lambda_\ell^n}{\rho^n} =: r(n) + w_n$$

where $r(n) \in O(n^{-1})$ is real-valued and the LRS $\langle w_n \rangle_n$ is both real-valued and simple. In addition, the characteristic roots of $\langle w_n \rangle_n$ are all non-real and lie on the unit circle. For the avoidance of doubt, the exponential polynomial defining $\langle w_n \rangle_n$ is real-valued since the summands $\alpha_\ell \lambda_\ell^n / \rho^n$ for non-real λ_ℓ occur in complex-conjugate pairs. Thus, w_n satisfies the hypothesis in Lemma 17, and so the inequalities $v_n < r(n) + w_n < r(n) + c$ hold for some $c < 0$ and infinitely many n . Since $r(n) \in O(n^{-1})$, we find that for infinitely many n , $v_n < 0$. Hence $\langle u_n \rangle_n$ is neither positive nor ultimately positive. \triangleleft

5 Hard Instances

In this section we discuss obstacles to extending our results for deciding positivity of reversible LRS of higher orders. Specifically, we construct a simple reversible LRS of order 18 and sketch the construction of a reversible LRS of order 12 that, to the best of our knowledge, lie outside the known classes for which the Positivity Problem can be decided. In particular, these examples lie beyond the scope of Theorem 4.

We start with simple reversible LRS of order 18. In order to illustrate the technical arguments and guide our construction of a hard instance, it is useful to recall the techniques employed by Ouaknine and Worrell in their proof of Theorem 4 [13]. For the sake of brevity, we shall give only a brief outline here; we direct the interested reader to the full argument given in [13].

5.1 Sketch proof of Theorem 4

Let $\langle u_n \rangle_n$ be a simple LRS satisfying the assumptions of Theorem 4. We first normalise $\langle u_n \rangle_n$ and so assume that the dominant roots $\lambda_1, \dots, \lambda_k$ of $\langle u_n \rangle_n$ lie on the unit circle in the complex plane. Then, for each $n \in \mathbb{N}$,

$$u_n = \alpha_1 \lambda_1^n + \dots + \alpha_k \lambda_k^n + \beta_1 \xi_1^n + \dots + \beta_{k'} \xi_{k'}^n$$

where $\xi_1, \dots, \xi_{k'}$ are the non-dominant roots of $\langle u_n \rangle_n$ and $\alpha_1, \dots, \alpha_k, \beta_1, \dots, \beta_{k'}$ are algebraic numbers.

We then compute a basis for the multiplicative relations between the dominant roots and consider a maximal subset $\lambda_1, \dots, \lambda_\ell$ whose elements are multiplicatively independent. By Kronecker’s Theorem on simultaneous Diophantine approximation (cf. [2, page 53]), $\{(\lambda_1^n, \dots, \lambda_\ell^n) : n \in \mathbb{N}\}$ is a dense subset of the torus $T := \{z \in \mathbb{C} : |z| = 1\}^\ell$, which is compact.

Ouaknine and Worrell then construct a continuous function $\tau : T \rightarrow \mathbb{R}$ given by

$$\tau(\lambda_1^n, \dots, \lambda_\ell^n) = \alpha_1 \lambda_1^n + \dots + \alpha_k \lambda_k^n$$

with the following properties. If $\min_T \tau = 0$, the given sequence $\langle u_n \rangle_n$ is ultimately positive. That is to say, there is a number N such that $u_n \geq 0$ for all $n \geq N$. If $\min_T \tau < 0$, the sequence is not ultimately positive (and thus also not positive). Finally, if $\min_T \tau > 0$, then the sequence grows quickly, and deciding positivity is relatively straightforward. Hence the critical case occurs when $\min_T \tau = 0$. Moreover, we can determine which of the three cases occur (that is, compute $\min_T \tau$).

In the critical case where $\min_T \tau = 0$, we can sometimes exploit the set of points $Z = \{(z_1, \dots, z_\ell) \in T : \tau(z_1, \dots, z_\ell) = 0\}$ where the minimum is attained. If $(z_1, \dots, z_\ell) \in Z$, then Baker’s Theorem on linear forms shows that λ_1^n cannot get too “close” to z_1 for n greater than a computable bound. As such, if Z is finite, then we can decide whether $\langle u_n \rangle_n$ is positive.

Theorem 4 is now proven as follows. If $\langle u_n \rangle_n$ has at most eight dominant characteristic roots and falls into the critical case, then Z is finite. Likewise, if $\langle u_n \rangle_n$ has exactly nine characteristic roots all of which are dominant, then $\langle u_n \rangle_n$ is positive in the critical case as $u_n \geq \min_T \tau = 0$.

The approach described breaks down when there are nine dominant roots since then Z is possibly infinite. Briefly, in this setting the state of the art cannot show that $(\lambda_1^n, \dots, \lambda_\ell^n)$ does not approach this infinite set too “closely”. Thus we encounter examples of LRS where we cannot currently determine positivity.

5.2 Constructing a hard example of a simple sequence of order 18

Our hard example is constructed from a function τ that assumes its minimum infinitely often on the torus $T = \{(z_1, z_2) \in \mathbb{C} : |z_1| = |z_2| = 1\}$. To this end, we define $\tau : T \rightarrow \mathbb{R}$ by

$$\tau(z_1, z_2) = (az_1 + \bar{a}z_1^{-1} + bz_2 + \bar{b}z_2^{-1})^2 \tag{5}$$

for some non-zero $a, b \in \mathbb{C}$ with $|a| \neq |b|$. Then $\min_T \tau$ is equal to 0 and τ attains its minimum on an infinite subset of T . This property prevents the application of Theorem 4.

► **Example 18.** We shall construct a simple reversible LRS sequence of order 18. An analysis of the spectral properties of this sequence shows that it lies beyond the current state-of-the-art techniques for deciding positivity. This hard instance is derived from the irreducible polynomial

$$f(X) = X^8 - 3X^7 + 4X^6 - 4X^5 + 11X^4 - 21X^3 + 19X^2 - 7X + 1,$$

130:14 Positivity Problems for Reversible Linear Recurrence Sequences

which has eight non-real roots $\lambda_1, \dots, \bar{\lambda}_4$ such that λ_1 and λ_2 are dominant, λ_3 and λ_4 are both non-dominant, and $1.143 \approx |\lambda_3| > 1 > |\lambda_4|$.

Let $\phi := (1 + \sqrt{5})/2$ denote the golden ratio. Then, with a certain labelling of complex conjugates,

$$\lambda_1 \bar{\lambda}_1 = \lambda_2 \bar{\lambda}_2 = \phi^2 \quad \text{and} \quad \lambda_3 \lambda_4 = \bar{\lambda}_3 \bar{\lambda}_4 = \phi^{-2},$$

which, due to the number of relations, severely limits the possible Galois automorphisms. In particular, the Galois group has the form of a *wreath product* $D_4 \wr C_2$. Thus a dihedral group D_4 acts on $\lambda_1, \bar{\lambda}_1, \lambda_2$, and $\bar{\lambda}_2$ and is generated by the elements (written in cycle notation) $(\lambda_1 \lambda_2 \bar{\lambda}_1 \bar{\lambda}_2)$ and $(\lambda_1 \bar{\lambda}_1)$. A second dihedral group D_4 acts on $\lambda_3, \bar{\lambda}_3, \lambda_4, \bar{\lambda}_4$ and is generated by $(\lambda_3 \bar{\lambda}_3 \lambda_4 \bar{\lambda}_4)$ and $(\lambda_3 \lambda_4)$. Lastly, there is a cyclic C_2 group acting on these two sets of four roots generated by the permutation $(\lambda_1 \lambda_3)(\bar{\lambda}_1 \bar{\lambda}_3)(\lambda_2 \bar{\lambda}_2)(\bar{\lambda}_2 \bar{\lambda}_4)$.

The terms in the sequence $\langle u_n \rangle_n$ are given as follows:

$$u_n = \frac{1}{\sqrt{5}} \left((1 + \lambda_1) \lambda_1^n + (1 + \bar{\lambda}_1) \bar{\lambda}_1^n + (1 + \lambda_2) \lambda_2^n + (1 + \bar{\lambda}_2) \bar{\lambda}_2^n \right)^2 \\ - \frac{1}{\sqrt{5}} \left((1 + \lambda_3) \lambda_3^n + (1 + \bar{\lambda}_3) \bar{\lambda}_3^n + (1 + \lambda_4) \lambda_4^n + (1 + \bar{\lambda}_4) \bar{\lambda}_4^n \right)^2.$$

By the action of the Galois group, it can be seen that each term u_n is rational and further that $\langle u_n \rangle_n$ is simple, reversible, and has exactly order 18. The initial values u_0, \dots, u_{17} of $\langle u_n \rangle_n$ are

$$-11, -8, 0, 240, 704, -20, 192, 5508, 46305, 2625, 13425, 73117, \\ 2469800, 536000, 554151, 77287, 108792361, 66461616.$$

The simple LRS $\langle u_n \rangle_n$ satisfies the relation

$$u_{n+18} = u_{n+17} - 10u_{n+16} + 6u_{n+15} + 43u_{n+14} - 93u_{n+13} + 672u_{n+12} - 596u_{n+11} \\ + 120u_{n+10} + 3972u_{n+9} - 15345u_{n+8} + 29654u_{n+7} - 36108u_{n+6} + 23847u_{n+5} \\ - 9572u_{n+4} + 2361u_{n+3} - 325u_{n+2} + 26u_{n+1} - u_n.$$

Observe that u_0, u_1 , and u_5 are negative, but up to $n = 10^5$ these are the only negative terms. Thus, the question is to prove that $u_n \geq 0$ for all $n \geq 6$. We reiterate that, as far as the authors are aware, there are no known techniques in the state of the art that can tackle this question.

It remains to show that the torus T associated with $\langle u_n \rangle_n$ has the prescribed “squaring form” (as in (5)) and that $\langle u_n \rangle_n$ is non-degenerate. To start, u_n is positive if and only if $\frac{u_n}{\phi^{2n}}$. Moreover, we observe that $|1 + \lambda_1| \neq |1 + \lambda_2|$ and that both λ_1/ϕ and λ_2/ϕ lie on the unit circle. For $a = 1 + \lambda_1, b = \lambda_2$ and some $0 < r < 1$, we have that

$$\frac{u_n}{\phi^{2n}} = \frac{1}{\phi^{2n}} \left((1 + \lambda_1) \lambda_1^n + (1 + \bar{\lambda}_1) \bar{\lambda}_1^n + (1 + \lambda_2) \lambda_2^n + (1 + \bar{\lambda}_2) \bar{\lambda}_2^n \right)^2 + O(r^n) \\ = \left(a \left(\frac{\lambda_1}{\phi} \right)^n + \bar{a} \left(\frac{\lambda_1}{\phi} \right)^{-n} + b \left(\frac{\lambda_2}{\phi} \right)^n + \bar{b} \left(\frac{\lambda_2}{\phi} \right)^{-n} \right)^2 + O(r^n)$$

is close to the “squaring form” discussed at (5). In fact, we have that

$$u_n/\phi^{2n} = \tau((\lambda_1/\phi)^n, (\lambda_2/\phi)^n) + O(r^n).$$

Here, the term $O(r^n)$ decreases exponentially fast and determines how closely the square should approach zero to contradict positivity.

We now show that we cannot apply Theorem 4 in this instance. To this end, we need to show that the points to which we restrict τ are dense on the torus T . That is, we need to show that λ_1/ϕ and λ_2/ϕ are multiplicatively independent. This lack of multiplicative relations also immediately implies that $\langle u_n \rangle_n$ is non-degenerate. We complete the spectral analysis of sequence $\langle u_n \rangle_n$ with the following proposition.

► **Proposition 19.** *We have that $\lambda_1/|\lambda_1|$ and $\lambda_2/|\lambda_2|$ are multiplicatively independent.*

Proof. Note that $|\lambda_1| = |\lambda_2| = \phi$ as $\lambda_1\bar{\lambda}_1 = \lambda_2\bar{\lambda}_2 = \phi^2$. By the earlier described Galois action, we see that there are Galois automorphisms σ and τ such that $\sigma(\lambda_1) = \tau(\lambda_1) = \lambda_3$, $\sigma(\lambda_2) = \lambda_4$ and $\tau(\lambda_2) = \bar{\lambda}_3$. Further, by this choice, $\sigma(\phi) = \tau(\phi) = -\phi^{-1}$.

Assume, for a contradiction, that $\lambda_1/|\lambda_1|$ and $\lambda_2/|\lambda_2|$ are multiplicatively dependent; that is to say, there are $a, b \in \mathbb{Z}$, not both 0, such that $(\lambda_1/|\lambda_1|)^a (\lambda_2/|\lambda_2|)^b = 1$. By applying σ to this identity we obtain

$$1 = \left(\frac{\lambda_3}{-\phi^{-1}}\right)^a \left(\frac{\lambda_4}{-\phi^{-1}}\right)^b = \zeta \left(\frac{|\lambda_3\lambda_4|}{\phi^{-2}}\right)^a \left(\frac{\lambda_4}{-\phi^{-1}}\right)^{b-a} = \zeta \left(\frac{\lambda_4}{-\phi^{-1}}\right)^{b-a}$$

for some ζ on the unit circle. Since $|\lambda_4/-\phi^{-1}| \neq 1$, we conclude that $a = b$. Then when we apply τ to the identity $(\lambda_1/|\lambda_1|)^a (\lambda_2/|\lambda_2|)^b = 1$ we obtain

$$1 = \left(\frac{\lambda_3}{-\phi^{-1}}\right)^a \left(\frac{\bar{\lambda}_3}{-\phi^{-1}}\right)^b = \zeta' \left(\frac{|\lambda_3|}{|\lambda_3|}\right)^a \left(\frac{\bar{\lambda}_3}{-\phi^{-1}}\right)^{b+a} = \zeta' \left(\frac{\bar{\lambda}_3}{-\phi^{-1}}\right)^{b+a}$$

for some ζ' on the unit circle. Since $|\bar{\lambda}_3/-\phi^{-1}| \neq 1$, this implies that $a = -b$. Together with $a = b$, we deduce that $a = b = 0$. Thus $\lambda_1/|\lambda_1|$ and $\lambda_2/|\lambda_2|$ are multiplicatively independent. ◀

5.3 Constructing a hard example of a non-simple sequence of order 12

In this subsection, we briefly consider a reversible LRS of order 12 where we cannot decide positivity nor ultimate positivity. Explicit examples are easier to construct than in the simple case and are closely related to the extensive discussion in [16]. Let us recall the following point from Theorem 1: a non-simple LRS that is a hard example of (ultimate) positivity possesses three simple dominant roots of which one is real and positive. One choice, closely resembling Example 4.5 in [11], is to take

$$\rho = \sqrt{2} + 1 \quad \text{and} \quad \lambda = \frac{1 + \sqrt{1 - 4\rho^2}}{2}.$$

Then we have that ρ and λ are units of equal modulus, ρ has one Galois conjugate $\tilde{\rho}$ of smaller modulus, and λ has three Galois conjugates. The three Galois conjugates of λ are its complex conjugate and two real numbers, say, λ_3 and λ_4 of smaller modulus. Lastly, let $q \in \mathbb{Q}_{>0}$. Then define the non-simple reversible rational-valued LRS $\langle u_n^q \rangle_n$ as follows:

$$u_n^q = (n + \rho)\rho^n + (n + \tilde{\rho})\tilde{\rho}^n + q(n + \lambda)\lambda^n + q(n + \bar{\lambda})\bar{\lambda}^n + q(n + \lambda_3)\lambda_3^n + q(n + \lambda_4)\lambda_4^n.$$

For small q , $\langle u_n^q \rangle_n$ is positive and so ultimately positive. For sufficiently large q , $\langle u_n^q \rangle_n$ is neither positive nor ultimately positive. However, given the current state of the art, it is not known how to determine where an arbitrary q falls in this partition. Thus, at the time of writing, we cannot tell whether LRS of the form $\langle u_n^q \rangle_n$ are (ultimately) positive.

Following [11, Section 4.2]), we can construct further LRS (akin to $\langle u_n^q \rangle_n$) where the state of the art is unable to settle positivity and ultimate positivity. In this direction, we may take a real quadratic unit $\rho > 1$ and find a non-real algebraic unit λ of equal modulus such that λ has a minimum polynomial of degree 4.

References

- 1 Mark Braverman. Termination of integer linear programs. In *International conference on computer aided verification*, pages 372–385. Springer, 2006.
- 2 J. W. S. Cassels. *An introduction to Diophantine approximation*. Cambridge Tracts in Mathematics and Mathematical Physics, No. 45. Cambridge University Press, New York, 1957.
- 3 Henri Cohen. *A course in computational algebraic number theory*, volume 138 of *Graduate Texts in Mathematics*. Springer-Verlag, Berlin, 1993.
- 4 The SageMath Developers. *SageMath, the Sage Mathematics Software System (Version 9.7)*, 2022. <https://www.sagemath.org>.
- 5 A. Dubickas and C. J. Smyth. On the Remak height, the Mahler measure and conjugate sets of algebraic numbers lying on two circles. *Proc. Edinb. Math. Soc. (2)*, 44(1):1–17, 2001. doi:10.1017/S001309159900098X.
- 6 Graham Everest, Alf van der Poorten, Igor Shparlinski, and Thomas Ward. *Recurrence sequences*, volume 104 of *Mathematical Surveys and Monographs*. Amer. Math. Soc., Providence, RI, 2003.
- 7 Ronald Ferguson. Irreducible polynomials with many roots of equal modulus. *Acta Arith.*, 78(3):221–225, 1997. doi:10.4064/aa-78-3-221-225.
- 8 Vesa Halava, Tero Harju, Mika Hirvensalo, and Juhani Karhumäki. Skolem’s problem—on the border between decidability and undecidability. Technical report, Turku Centre for Computer Science, 2005.
- 9 George Kenison. On the Skolem Problem for Reversible Sequences. In Stefan Szeider, Robert Ganian, and Alexandra Silva, editors, *47th International Symposium on Mathematical Foundations of Computer Science (MFCS 2022)*, volume 241 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 61:1–61:15, Dagstuhl, Germany, 2022. Schloss Dagstuhl – Leibniz-Zentrum für Informatik. doi:10.4230/LIPIcs.MFCS.2022.61.
- 10 L. Kronecker. Zwei Sätze über Gleichungen mit ganzzahligen Coefficienten. *Journal für die reine und angewandte Mathematik (Crelles Journal)*, 1857(53):173–175, January 1857. doi:10.1515/crll.1857.53.173.
- 11 Richard Lipton, Florian Luca, Joris Nieuwveld, Joël Ouaknine, David Purser, and James Worrell. On the Skolem Problem and the Skolem Conjecture. In *Proceedings of the 37th Annual ACM/IEEE Symposium on Logic in Computer Science, LICS ’22*, New York, NY, USA, 2022. Association for Computing Machinery. doi:10.1145/3531130.3533328.
- 12 Maurice Mignotte, Tarlok Shorey, and Robert Tijdeman. The distance between terms of an algebraic recurrence sequence. *Journal für die Reine und Angewandte Mathematik*, pages 63–76, 1984.
- 13 Joël Ouaknine and James Worrell. On the Positivity Problem for Simple Linear Recurrence sequences,. In Javier Esparza, Pierre Fraigniaud, Thore Husfeldt, and Elias Koutsoupias, editors, *Automata, Languages, and Programming*, pages 318–329, Berlin, Heidelberg, 2014. Springer Berlin Heidelberg.
- 14 Joël Ouaknine and James Worrell. Ultimate positivity is decidable for simple linear recurrence sequences. In Javier Esparza, Pierre Fraigniaud, Thore Husfeldt, and Elias Koutsoupias, editors, *Automata, Languages, and Programming - 41st International Colloquium, ICALP 2014, Copenhagen, Denmark, July 8-11, 2014, Proceedings, Part II*, volume 8573 of *Lecture Notes in Computer Science*, pages 330–341. Springer, 2014. doi:10.1007/978-3-662-43951-7_28.
- 15 Joël Ouaknine and James Worrell. On linear recurrence sequences and loop termination. *ACM SIGLOG News*, 2(2):4–13, April 2015.
- 16 Joël Ouaknine and James Worrell. *Positivity Problems for Low-Order Linear Recurrence Sequences*, pages 366–379. ACM, New York, 2014. doi:10.1137/1.9781611973402.27.
- 17 Alfred Pringsheim. Ueber Functionen, welche in gewissen Punkten endliche Differentialquotienten jeder endlichen Ordnung, aber keine Taylor’sche Reihenentwicklung besitzen. *Mathematische Annalen*, 44(1):41–56, 1894.

- 18 J.J. Rotman. *An Introduction to the Theory of Groups*. Graduate Texts in Mathematics. Springer New York, 2012.
- 19 C. Smyth. Conjugate algebraic numbers on conics. *Acta Arithmetica*, 40(4):333–346, 1982.
- 20 I. Stewart and D. Tall. *Algebraic number theory and Fermat’s last theorem*. CRC Press, Boca Raton, FL, fourth edition, 2016.
- 21 Edward Charles Titchmarsh. *The theory of functions*. Oxford University Press, 2nd edition, 1939.
- 22 Nikolai Vereshchagin. Occurrence of zero in a linear recursive sequence. *Mathematical notes of the Academy of Sciences of the USSR*, 38(2):609–615, August 1985.
- 23 Giulio Vivanti. Sulle serie di potenze. *Annali di Matematica Pura ed Applicata (1867-1897)*, 21(1):193–194, 1893.