

Semi-algebraic Ramsey Numbers

Andrew Suk

University of Illinois at Chicago
851 S. Morgan St., Chicago, IL 60607, USA
suk@uic.edu

Abstract

Given a finite point set $P \subset \mathbb{R}^d$, a k -ary semi-algebraic relation E on P is the set of k -tuples of points in P , which is determined by a finite number of polynomial equations and inequalities in kd real variables. The description complexity of such a relation is at most t if the number of polynomials and their degrees are all bounded by t . The Ramsey number $R_k^{d,t}(s, n)$ is the minimum N such that any N -element point set P in \mathbb{R}^d equipped with a k -ary semi-algebraic relation E such that E has complexity at most t , contains s members such that every k -tuple induced by them is in E or n members such that every k -tuple induced by them is not in E .

We give a new upper bound for $R_k^{d,t}(s, n)$ for $k \geq 3$ and s fixed. In particular, we show that for fixed integers d, t, s

$$R_3^{d,t}(s, n) \leq 2^{n^{o(1)}},$$

establishing a subexponential upper bound on $R_3^{d,t}(s, n)$. This improves the previous bound of $2^{n^{C_1}}$ due to Conlon, Fox, Pach, Sudakov, and Suk where C_1 depends on d and t , and improves upon the trivial bound of $2^{n^{C_2}}$ which can be obtained by applying classical Ramsey numbers where C_2 depends on s . As an application, we give new estimates for a recently studied Ramsey-type problem on hyperplane arrangements in \mathbb{R}^d . We also study multi-color Ramsey numbers for triangles in our semi-algebraic setting, achieving some partial results.

1998 ACM Subject Classification G.2.2 Graph Theory

Keywords and phrases Ramsey theory, semi-algebraic relation, one-sided hyperplanes, Schur numbers

Digital Object Identifier 10.4230/LIPIcs.SOCG.2015.59

1 Introduction

Classical Ramsey numbers. A k -uniform hypergraph $H = (P, E)$ consists of a vertex set P and an edge set $E \subset \binom{P}{k}$, which is a collection of subsets of P of size k . The Ramsey number $R_k(s, n)$ is the minimum integer N such that every k -uniform hypergraph on N vertices contains either s vertices such that every k -tuple induced by them is an edge, or contains n vertices such that every k -tuple induced by them is not an edge.

Due to its wide range of applications in logic, number theory, analysis, and geometry, estimating Ramsey numbers has become one of the most central problems in combinatorics. For *diagonal* Ramsey numbers, i.e. when $s = n$, the best known lower and upper bounds for $R_k(n, n)$ are of the form¹ $R_2(n, n) = 2^{\Theta(n)}$, and for $k \geq 3$,

$$\text{twr}_{k-1}(\Omega(n^2)) \leq R_k(n, n) \leq \text{twr}_k(O(n)),$$

¹ We write $f(n) = O(g(n))$ if $|f(n)| \leq c|g(n)|$ for some fixed constant c and for all $n \geq 1$; $f(n) = \Omega(g(n))$ if $g(n) = O(f(n))$; and $f(n) = \Theta(g(n))$ if both $f(n) = O(g(n))$ and $f(n) = \Omega(g(n))$ hold. We write $f(n) = o(g(n))$ if for every positive $\epsilon > 0$ there exists a constant n_0 such that $f(n) \leq \epsilon|g(n)|$ for all $n \geq n_0$.



© Andrew Suk;

licensed under Creative Commons License CC-BY

31st International Symposium on Computational Geometry (SoCG'15).

Editors: Lars Arge and János Pach; pp. 59–73



Leibniz International Proceedings in Informatics

LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

where the tower function $\text{twr}_k(x)$ is defined by $\text{twr}_1(x) = x$ and $\text{twr}_{i+1} = 2^{\text{twr}_i(x)}$ (see [21, 18, 19, 20]). Erdős, Hajnal, and Rado [19] conjectured that $R_k(n, n) = \text{twr}_k(\Theta(n))$, and Erdős offered a \$500 reward for a proof. Despite much attention over the last 50 years, the exponential gap between the lower and upper bounds for $R_k(n, n)$, when $k \geq 3$, remains unchanged.

The *off-diagonal* Ramsey numbers, i.e. $R_k(s, n)$ with s fixed and n tending to infinity, has also been extensively studied. Unlike $R_k(n, n)$, the lower and upper bounds for $R_k(s, n)$ are much more comparable. It is known [5, 25, 8, 9] that $R_2(3, n) = \Theta(n^2/\log n)$ and, for fixed $s > 3$

$$\Omega\left(n^{\frac{s+1}{2}-\epsilon}\right) \leq R_2(s, n) \leq O\left(n^{s-1}\right), \quad (1)$$

where $\epsilon > 0$ is an arbitrarily small constant. Combining the upper bound in (1) with the results of Erdős, Hajnal, and Rado [20, 19] demonstrates that

$$\text{twr}_{k-1}(\Omega(n)) \leq R_k(s, n) \leq \text{twr}_{k-1}(O(n^{2s-4})), \quad (2)$$

for $k \geq 3$ and $s \geq 2^k$. See Conlon, Fox, and Sudakov [14] for a recent improvement.

Semi-algebraic setting. In this paper, we continue a sequence of recent works on Ramsey numbers for k -ary semi-algebraic relations E on \mathbb{R}^d (see [10, 17, 13, 33]). Before we give its precise definition, let us recall two classic Ramsey-type theorems of Erdős and Szekeres.

► **Theorem 1** ([21]). *For $N = (s-1)(n-1) + 1$, let $P = (p_1, \dots, p_N) \subset \mathbb{R}$ be a sequence of N distinct real numbers. Then P contains either an increasing subsequence of length s , or a decreasing subsequence on length n .*

In fact, there are now at least 6 different proofs of Theorem 1 (see [32]). The other well-known result from [21] is the following theorem, which is often referred to as the Erdős-Szekeres cups-caps Theorem. Let X be a finite point set in the plane in general position.² We say that $X = (p_{i_1}, \dots, p_{i_s})$ forms an s -cup (s -cap) if X is in convex position and its convex hull is bounded above (below) by a single edge.

► **Theorem 2** ([21]). *For $N = \binom{n+s-4}{s-2} + 1$, let $P = (p_1, \dots, p_N)$ be a sequence of N points in the plane in general position. Then P contains either an s -cup or an n -cap.*

Theorems 1 and 2 can be generalized using the following semi-algebraic framework. Let $P = \{p_1, \dots, p_N\}$ be a sequence of N points in \mathbb{R}^d . Then we say that $E \subset \binom{P}{k}$ is a *semi-algebraic relation* on P with *complexity* at most t , if there are t polynomials $f_1, \dots, f_t \in \mathbb{R}[x_1, \dots, x_{kd}]$ of degree at most t , and a Boolean function Φ such that for $1 \leq i_1 < \dots < i_k \leq N$,

$$(p_{i_1}, \dots, p_{i_k}) \in E \quad \Leftrightarrow \quad \Phi(f_1(p_{i_1}, \dots, p_{i_k}) \geq 0, \dots, f_t(p_{i_1}, \dots, p_{i_k}) \geq 0) = 1.$$

We say that the relation $E \subset \binom{P}{k}$ is *symmetric* if $(p_{i_1}, \dots, p_{i_k}) \in E$ iff for all permutation π ,

$$\Phi(f_1(p_{\pi(i_1)}, \dots, p_{\pi(i_k)}) \geq 0, \dots, f_t(p_{\pi(i_1)}, \dots, p_{\pi(i_k)}) \geq 0) = 1.$$

Point sets $P \subset \mathbb{R}^d$ equipped with a k -ary semi-algebraic relation $E \subset \binom{P}{k}$ are often used to model problems in discrete geometry, where the dimension d , uniformity k , and complexity

² No two members share the same x -coordinate, and no three members are collinear.

t are considered fixed but arbitrarily large constants. Since we can always make any relation E symmetric by increasing its complexity to $t' = t'(k, d, t)$, we can therefore simplify our presentation by only considering symmetric relations.

Let $R_k^{d,t}(s, n)$ be the minimum integer N such that every N -element point set P in \mathbb{R}^d equipped with a k -ary (symmetric) semi-algebraic relation $E \subset \binom{P}{k}$, which has complexity at most t , contains s points such that every k -tuple induced by them is in E , or contains n points such that no k -tuple induced by them is in E . Alon, Pach, Pinchasi, Radoičić, and Sharir [6] showed that for $k = 2$, we have

$$R_2^{d,t}(n, n) \leq n^C, \quad (3)$$

where $C = C(d, t)$. Roughly speaking, $C \approx t \binom{d+t}{t}$. Conlon, Fox, Pach, Sudakov, and Suk showed that one can adapt the Erdős-Rado argument in [20] and establish the following recursive formula for $R_k^{d,t}(s, n)$.

► **Theorem 3** ([13]). *Set $M = R_{k-1}^{d,t}(s-1, n-1)$. Then for every $k \geq 3$,*

$$R_k^{d,t}(s, n) \leq 2^{C_1 M \log M},$$

where $C_1 = C_1(k, d, t)$.

Together with (3) we have $R_k^{d,t}(n, n) \leq \text{twr}_{k-1}(n^C)$, giving an exponential improvement over the Ramsey numbers for general k -uniform hypergraphs. Conlon et al. [13] also gave a construction of a geometric example that provides a $\text{twr}_{k-1}(\Omega(n))$ lower bound, demonstrating that $R_k^{d,t}(n, n)$ does indeed grow as a $(k-1)$ -fold exponential tower in n .

However, off-diagonal Ramsey numbers for semi-algebraic relations are much less understood. The best known upper bound for $R_k^{d,t}(s, n)$ is essentially the trivial bound

$$R_k^{d,t}(s, n) \leq \min \left\{ R_k^{d,t}(n, n), R_k(s, n) \right\}.$$

The crucial case is when $k = 3$, since any significant improvement on estimating $R_3^{d,t}(s, n)$ could be used in combination with Theorem 3 to obtain a better bound for $R_k^{d,t}(s, n)$, for $k \geq 4$. The trivial bound implies that

$$R_3^{d,t}(s, n) \leq 2^{n^C}, \quad (4)$$

where C is a large constant depending on d, t , and s .

The main difficulty in improving (4) is that the Erdős-Rado upper bound argument [20] will not be effective. Roughly speaking, the Erdős-Rado argument reduces the problem from 3-uniform hypergraphs to graphs, producing a recursive formula similar to Theorem 3. This approach has been used repeatedly by many researchers to give upper bounds on Ramsey-type problems arising in triple systems [14, 13, 33, 30]. However, it is very unlikely that any variant of the Erdős-Rados upper bound argument will establish a subexponential upper bound for $R_3^{d,t}(s, n)$.

With a more novel approach, our main result establishes the following improved upper bound for $R_3^{d,t}(s, n)$, showing that the function $R_3^{d,t}(s, n)$ is indeed subexponential in n .

► **Theorem 4.** *For fixed integers $d, t \geq 1$ and $s \geq 4$, we have $R_3^{d,t}(s, n) \leq 2^{n^{o(1)}}$. More precisely,*

$$R_3^{d,t}(s, n) \leq 2^{2^{c\sqrt{(\log n)(\log \log n)}}},$$

where $c = c(d, t, s)$.

Combining Theorems 4 and 3 we have the following.

► **Corollary 5.** *For fixed integers $d, t \geq 1$, $k \geq 3$, and $s \geq k + 1$, we have*

$$R_k^{d,t}(s, n) \leq \text{twr}_{k-1}(n^{o(1)}).$$

For $d \geq 2$ and $t \geq 1$, the classic cups-caps construction of Erdős and Szekeres [21] shows that $R_3^{d,t}(s, n) \geq \Omega(n^{s-2})$, and together with the semi-algebraic stepping-up lemma proven in [13] (see also [16]) we have $R_k^{d,t}(s, n) \geq \text{twr}_{k-2}(\Omega(n^{s-2}))$ for $s, d \geq 2^k$.

In Section 5, we give an application of Theorem 4 to a recently studied problem on hyperplane arrangements in \mathbb{R}^d .

Monochromatic triangles. Let $R_2(s; m) = R_2(\underbrace{s, \dots, s}_m)$ denote the smallest integer N such that any m -coloring on the edges of the complete N -vertex graph contains a monochromatic clique of size s , that is, a set of s vertices such that every pair from this set has the same color. For the case $s = 3$, the Ramsey number $R_2(3; m)$ has received a lot of attention over the last 100 years due to its application in additive number theory [31] (more details are given in Section 6.1). It is known (see [24, 31]) that

$$\Omega(3.19^m) \leq R_2(3; m) \leq O(m!).$$

Our next result states that we can improve the upper bound on $R_2(3; m)$ in our semi-algebraic setting. More precisely, let $R_2^{d,t}(3; m)$ be the minimum integer N such that every N -element point set P in \mathbb{R}^d equipped with symmetric semi-algebraic relations $E_1, \dots, E_m \subset \binom{P}{2}$, such that each E_i has complexity at most t and $\binom{P}{2} = E_1 \cup \dots \cup E_m$, contains three points such that every pair induced by them belongs to E_i for some fixed i .

► **Theorem 6.** *For fixed $d, t \geq 1$ we have*

$$R_2^{d,t}(3; m) < 2^{O(m \log \log m)}.$$

We also show that for fixed $d \geq 1$ and $t \geq 5000$, the function $R_2^{d,t}(3; m)$ does indeed grow exponentially in m .

► **Theorem 7.** *For $d \geq 1$ and $t \geq 5000$ we have*

$$R_2^{d,t}(3; m) \geq c(1681)^{m/7} \geq c(2.889)^m,$$

where c is an absolute constant.

Organization. In the next two sections, we recall several old theorems on the arrangement of surfaces in \mathbb{R}^d and establish a result on point sets equipped with multiple binary relations. In Section 4, we combine the results from Sections 2 and 3 to prove our main result, Theorem 4. We discuss a short proof of our application in Section 5, and our results on monochromatic triangles in Section 6.

We systemically omit floor and ceiling signs whenever they are not crucial for the sake of clarity of our presentation. All logarithms are assumed to be base 2.

2 Arrangement of surfaces in \mathbb{R}^d

In this section, we recall several old results on the arrangement of surfaces in \mathbb{R}^d . Let f_1, \dots, f_m be d -variate real polynomials of degree at most t , with zero sets Z_1, \dots, Z_m , that is, $Z_i = \{x \in \mathbb{R}^d : f_i(x) = 0\}$. Set $\Sigma = \{Z_1, \dots, Z_m\}$. We will assume that d and t are fixed, and m is some number tending to infinity. A *cell* in the arrangement $\mathcal{A}(\Sigma) = \bigcup_i Z_i$ is a relatively open connected set defined as follows. Let \approx be an equivalence relation on \mathbb{R}^d , where $x \approx y$ if $\{i : x \in Z_i\} = \{i : y \in Z_i\}$. Then the cells of the arrangement $\mathcal{A}(\Sigma)$ are the connected components of the equivalence classes. A vector $\sigma \in \{-1, 0, +1\}^m$ is a *sign pattern* of f_1, \dots, f_m if there exists an $x \in \mathbb{R}^d$ such that the sign of $f_j(x)$ is σ_j for all $j = 1, \dots, m$. The Milnor-Thom theorem (see [7, 29, 34]) bounds the number of cells in the arrangement of the zero sets Z_1, \dots, Z_m and, consequently, the number of possible sign patterns (see all [35]).

► **Theorem 8** (Milnor-Thom). *Let f_1, \dots, f_m be d -variate real polynomials of degree at most t . The number of cells in the arrangement of their zero sets $Z_1, \dots, Z_m \subset \mathbb{R}^d$ and, consequently, the number of sign patterns of f_1, \dots, f_m is at most*

$$\left(\frac{50mt}{d}\right)^d,$$

for $m \geq d \geq 1$.

While the Milnor-Thom Theorem bounds the number of cells in the arrangement $\mathcal{A}(\Sigma)$, the complexity of these cells may be very large (depending on m). A long standing open problem is whether each cell can be further decomposed into semi-algebraic sets³ with bounded description complexity (which depends only on d and t), such that the total number of cells for the whole arrangement is still $O(m^d)$. This can be done easily in dimension 2 by a result of Chazelle et al. [11]. Unfortunately in higher dimensions, the current bounds for this problem are not tight. In dimension 3, Chazelle et al. [11] established a near tight bound of $O(m^3 \beta(m))$, where $\beta(m)$ is an extremal slowly growing function of m related to the inverse Ackermann function. For dimensions $d \geq 4$, Koltun [26] established a general bound of $O(m^{2d-4+\epsilon})$ for arbitrarily small constant ϵ , which is nearly tight in dimension 4. By combining these bounds with the standard theory of random sampling [4, 12, 6], one can obtain the following result which is often referred to as the Cutting Lemma. We say that the surface $Z_i = \{x \in \mathbb{R}^d : f_i(x) = 0\}$ *crosses* the cell $\Delta \subset \mathbb{R}^d$ if $Z_i \cap \Delta \neq \emptyset$ and Z_i does not fully contain Δ .

► **Lemma 9** (Cutting Lemma). *For $d, t \geq 1$, let Σ be a family of m algebraic surfaces (zero sets) in \mathbb{R}^d of degree at most t . Then for any integer $r \geq 1$, there exists a decomposition of \mathbb{R}^d into at most $c_1 r^{2d}$ relatively open connected sets (cells), where $c_1 = c_1(d, t)$, such that each cell is crossed by at most m/r surfaces from Σ .*

As an application, we prove the following lemma (see [27, 3] for a similar result when Σ is a collection of hyperplanes).

► **Lemma 10.** *For $d, t \geq 1$, let P be an N -element point set in \mathbb{R}^d and let Σ be a family of m surfaces of degree at most t . Then for any integer $\ell > \log m$, we can find ℓ disjoint subsets*

³ A real semi-algebraic set in \mathbb{R}^d is the locus of all points that satisfy a given finite Boolean combination of polynomial equations and inequalities in the d coordinates.

P_i and ℓ cells Δ_i , with $\Delta_i \supset P_i$, such that each subset P_i contains at least $N/(4\ell)$ points from P , and every surface in Σ crosses at most $c_2\ell^{1-1/(2d)}$ cells Δ_i , where $c_2 = c_2(d, t)$.

Proof. We first find Δ_1 and P_1 as follows. Let $\ell > \log m$ and let c_1 be as defined in Lemma 9. Given a family Σ of m surfaces in \mathbb{R}^d , we apply Lemma 9 with parameter $r = (\ell/c_1)^{1/(2d)}$, and decompose \mathbb{R}^d into at most ℓ cells, such that each cell is crossed by at most $\frac{m}{(\ell/c_1)^{1/(2d)}}$ surfaces from Σ . By the pigeonhole principle, there is a cell Δ_1 that contains at least N/ℓ points from P . Let P_1 be a subset of exactly $\lfloor N/\ell \rfloor$ points in $\Delta_1 \cap P$. Now for each surface from Σ that crosses Δ_1 , we "double it" by adding another copy of that surface to our collection. This gives us a new family of surfaces Σ_1 such that

$$|\Sigma_1| \leq m + \frac{m}{(\ell/c_1)^{1/(2d)}} = m \left(1 + \frac{1}{(\ell/c_1)^{1/(2d)}} \right).$$

After obtaining subsets P_1, \dots, P_i such that $|P_j| = \lfloor \frac{N}{\ell}(1 - \frac{1}{\ell})^{j-1} \rfloor$ for $1 \leq j \leq i$, cells $\Delta_1, \dots, \Delta_i$, and the family of surfaces Σ_i such that

$$|\Sigma_i| \leq m \left(1 + \frac{1}{(\ell/c_1)^{1/(2d)}} \right)^i,$$

we obtain $P_{i+1}, \Delta_{i+1}, \Sigma_{i+1}$ as follows. Given Σ_i , we apply Lemma 9 with the same parameter $r = (\ell/c_1)^{1/(2d)}$, and decompose \mathbb{R}^d into at most ℓ cells, such that each cell is crossed by at most $\frac{|\Sigma_i|}{(\ell/c_1)^{1/(2d)}}$ surfaces from Σ_i . Let $P' = P \setminus (P_1 \cup \dots \cup P_i)$. By the pigeonhole principle, there is a cell Δ_{i+1} that contains at least

$$\begin{aligned} \frac{|P'|}{\ell} &\geq \left(N - \sum_{j=1}^i \frac{N}{\ell} (1 - \frac{1}{\ell})^{j-1} \right) / \ell \\ &= \frac{N}{\ell} \left(1 - \frac{1}{\ell} \sum_{j=1}^i (1 - \frac{1}{\ell})^{j-1} \right) \\ &= \frac{N}{\ell} (1 - \frac{1}{\ell}) \left(1 - \frac{1}{\ell} - \frac{1}{\ell} \sum_{j=1}^{i-1} (1 - \frac{1}{\ell})^{j-1} \right) \\ &= \frac{N}{\ell} (1 - \frac{1}{\ell})^i \end{aligned}$$

points from P' . Let P_{i+1} be a subset of exactly $\lfloor \frac{N}{\ell} (1 - 1/\ell)^i \rfloor$ points in $\Delta_{i+1} \cap P'$. Finally, for each surface from Σ_i that crosses Δ_{i+1} , we "double it" by adding another copy of that surface to our collection, giving us a new family of surfaces Σ_{i+1} such that

$$\begin{aligned} |\Sigma_{i+1}| &\leq |\Sigma_i| + \frac{|\Sigma_i|}{(\ell/c_1)^{1/(2d)}} \\ &= |\Sigma_i| \left(1 + \frac{1}{(\ell/c_1)^{1/(2d)}} \right) \\ &\leq m \left(1 + \frac{1}{(\ell/c_1)^{1/(2d)}} \right)^{i+1}. \end{aligned}$$

Notice that $|P_i| \geq N/(4\ell)$ for $i \leq \ell$. Once we have obtained subsets P_1, \dots, P_ℓ and cell $\Delta_1, \dots, \Delta_\ell$, it is easy to see that each surface in Σ crosses at most $O(r^{1-1/(2d)})$ cells Δ_i . Indeed suppose $Z \in \Sigma$ crosses κ cells. Then by the arguments above, there must be 2^κ copies of Z in Σ_ℓ . Hence we have

$$2^\kappa \leq m \left(1 + \frac{1}{(\ell/c_1)^{1/(2d)}} \right)^\ell \leq m e^{c_1 \ell^{1-1/(2d)}}.$$

Since $\ell \geq \log m$, we have

$$\kappa \leq c_2 \ell^{1-1/(2d)},$$

for sufficiently large $c_2 = c_2(d, t)$. ◀

3 Multiple binary relations

Let P be a set of N points in \mathbb{R}^d , and let $E_1, \dots, E_m \subset \binom{P}{2}$ be binary (symmetric) semi-algebraic relations on P such that E_i has complexity at most t . The goal of this section is to find a large subset $P' \subset P$ such that $\binom{P'}{2} \cap E_i = \emptyset$ for all i , given that the clique number in the graphs $G_i = (P, E_i)$ are small.

First we recall a classic theorem of Dilworth (see also [23]). Let $G = (V, E)$ be a graph whose vertices are ordered $V = \{v_1, \dots, v_N\}$. We say that E is *transitive* on V if for $1 \leq i_1 < i_2 < i_3 \leq N$, $(v_{i_1}, v_{i_2}), (v_{i_2}, v_{i_3}) \in E$ implies that $(v_{i_1}, v_{i_3}) \in E$.

► **Theorem 11** (Dilworth). *Let $G = (V, E)$ be an N -vertex graph whose vertices are ordered $V = \{v_1, \dots, v_N\}$, such that E is transitive on V . If G has clique number ω , then G contains an independent set of size N/ω .*

► **Lemma 12.** *For integers $m \geq 2$ and $d, t \geq 1$, let P be a set of N points in \mathbb{R}^d equipped with (symmetric) semi-algebraic relations $E_1, \dots, E_m \subset \binom{P}{2}$, where each E_i has complexity at most t . Then there is a subset $P' \subset P$ of size $N^{1/(c_3 \log m)}$, where $c_3 = c_3(d, t)$, and a fixed ordering on P' such that each relation E_i is transitive on P' .*

Proof. We proceed by induction on N . Let c_3 be a sufficiently large number depending only on d and t that will be determined later. For each relation $E_i \subset \binom{P}{2}$, let $f_{i,1}, \dots, f_{i,t}$ be polynomials of degree at most t and let Φ_i be a boolean function such that

$$(p, q) \in E_i \iff \Phi_i(f_{i,1}(p, q) \geq 0, \dots, f_{i,t}(p, q) \geq 0) = 1.$$

For each $p \in P$, $i \in \{1, \dots, m\}$, and $j \in \{1, \dots, t\}$, we define the surface $Z_{p,i,j} = \{x \in \mathbb{R}^d : f_{i,j}(p, x) = 0\}$. Then let Σ be the family of Nmt surfaces in \mathbb{R}^d defined by

$$\Sigma = \{Z_{p,i,j} : p \in P, 1 \leq i \leq m, 1 \leq j \leq t\}.$$

By applying Lemma 9 to Σ with parameter $r = (mt)^2$, there is a decomposition of \mathbb{R}^d into at most $c_1(mt)^{4d}$ cells such that each cell has the property that at most $N/(mt)$ surfaces from Σ crosses it. We note that $c_1 = c_1(d, t)$ is defined in Lemma 9. By the pigeonhole principle, there is a cell Δ in the decomposition such that $|\Delta \cap P| \geq N/(c_1(mt)^{4d})$. Set $P_1 = \Delta \cap P$.

Let $P_2 \subset P \setminus P_1$ such that each point in P_2 gives rise to mt surfaces that do not cross Δ . More precisely,

$$P_2 = \{p \in P \setminus P_1 : Z_{p,i,j} \text{ does not cross } \Delta, \forall i, j\}.$$

Notice that

$$|P_2| \geq N - \frac{N}{mt} - \frac{N}{c_1(mt)^{4d}} \geq \frac{N}{4}.$$

We fix a point $p_0 \in P_1$. Then for each $q \in P_2$, let $\sigma(q) \in \{-1, 0, +1\}^{mt}$ be the sign pattern of the (mt) -tuple $(f_{1,1}(p_0, q), f_{1,2}(p_0, q), \dots, f_{m,t}(p_0, q))$. By Theorem 8, there are at most

$\left(\frac{50mt^2}{d}\right)^d$ distinct sign vectors σ . By the pigeonhole principle, there is a subset $P_3 \subset P_2$ such that

$$|P_3| \geq \frac{|P_2|}{(50/d)^d m^d t^{2d}},$$

and for any two points $q, q' \in P_3$, we have $\sigma(q) = \sigma(q')$. That is, q and q' give rise to vectors with the same sign pattern. Therefore, for any $p, p' \in P_1$ and $q, q' \in P_3$, we have $(p, q) \in E_i$ if and only if $(p', q') \in E_i$, for all $i \in \{1, \dots, m\}$.

Let $c_4 = c_4(d, t)$ be sufficiently large such that $|P_1|, |P_3| \geq \frac{N}{c_4 m^{4d}}$. By the induction hypothesis, we can find subsets $P_4 \subset P_1, P_5 \subset P_3$, such that

$$|P_4|, |P_5| \geq \left(\frac{N}{c_4 m^{4d}}\right)^{\frac{1}{c_3 \log m}} \geq \frac{N^{\frac{1}{c_3 \log m}}}{2},$$

where $c_3 = c_3(d, t)$ is sufficiently large, and there is an ordering on P_4 (and on P_5) such that each E_i is transitive on P_4 (and on P_5). Set $P' = P_4 \cup P_5$, which implies $|P'| \geq N^{\frac{1}{c_3 \log m}}$. We will show that P' has the desired properties. Let π and π' be the orderings on P_4 and P_5 respectively, such that E_i is transitive on P_4 and on P_5 , for every $i \in \{1, \dots, m\}$. We order the elements in $P' = \{p_1, \dots, p_{|P'|}\}$ by using π and π' , such that all elements in P_5 comes after all elements in P_4 .

In order to show that E_i is transitive on P' , it suffices to examine triples going across P_4 and P_5 . Let $p_{j_1}, p_{j_2} \in P_4$ and $p_{j_3} \in P_5$ such that $j_1 < j_2 < j_3$. By construction of P_4 and P_5 , if $(p_{j_1}, p_{j_2}), (p_{j_2}, p_{j_3}) \in E_i$, then we have $(p_{j_1}, p_{j_3}) \in E_i$. Likewise, suppose $p_{j_1} \in P_4$ and $p_{j_2}, p_{j_3} \in P_5$. Then again by construction of P_4 and P_5 , if $(p_{j_1}, p_{j_2}), (p_{j_2}, p_{j_3}) \in E_i$, then we have $(p_{j_1}, p_{j_3}) \in E_i$. Hence E_i is transitive on P' , for all $i \in \{1, \dots, m\}$, and this completes the proof. \blacktriangleleft

By combining the two previous results, we have the following.

► **Lemma 13.** *For $m \geq 2$ and $d, t \geq 1$, let P be a set of N points in \mathbb{R}^d equipped with (symmetric) semi-algebraic relations $E_1, \dots, E_m \subset \binom{P}{2}$, where each E_i has complexity at most t . If graph $G_i = (P, E_i)$ has clique number ω_i , then there is a subset $P' \subset P$ of size $\frac{N^{1/(c_3 \log m)}}{\omega_1 \cdots \omega_m}$, where $c_3 = c_3(d, t)$ is defined above, such that $\binom{P'}{2} \cap E_i = \emptyset$ for all i .*

Proof. By applying Lemma 12, we obtain a subset $P_1 \subset P$ of size $N^{\frac{1}{c_3 \log m}}$, and an ordering on P_1 such that E_i is transitive on P_1 for all i . Then by an m -fold application of Theorem 11, the statement follows. \blacktriangleleft

4 Proof of Theorem 4

Let P be a point set in \mathbb{R}^d and let $E \subset \binom{P}{3}$ be a semi-algebraic relation on P . We say that (P, E) is $K_s^{(3)}$ -free if every collection of s points in P contains a triple not in E . Suppose we have ℓ disjoint subsets $P_1, \dots, P_\ell \subset P$. For $1 \leq i_1 < i_2 < i_3 \leq \ell$, we say that the triple $(P_{i_1}, P_{i_2}, P_{i_3})$ is *homogeneous* if $(p_1, p_2, p_3) \in E$ for all $p_1 \in P_{i_1}, p_2 \in P_{i_2}, p_3 \in P_{i_3}$, or $(p_1, p_2, p_3) \notin E$ for all $p_1 \in P_{i_1}, p_2 \in P_{i_2}, p_3 \in P_{i_3}$. For $p_1, p_2 \in P_1 \cup \dots \cup P_\ell$ and $i \in \{1, \dots, \ell\}$, we say that the triple (p_1, p_2, i) is *good*, if $(p_1, p_2, p_3) \in E$ for all $p_3 \in P_i$, or $(p_1, p_2, p_3) \notin E$ for all $p_3 \in P_i$. We say that the triple (p_1, p_2, i) is *bad* if (p_1, p_2, i) is not good and $p_1, p_2 \notin P_i$.

► **Lemma 14.** *Let P be a set of N points in \mathbb{R}^d and let $E \subset \binom{P}{3}$ be a (symmetric) semi-algebraic relation on P such that E has complexity at most t . Then for $r = \frac{N^{1/(30d)}}{tc_2}$, where c_2 is defined in Lemma 10, there are disjoint subsets $P_1, \dots, P_r \subset P$ such that*

1. $|P_i| \geq \frac{N^{1/(30d)}}{tc_2}$,
2. all triples $(P_{i_1}, P_{i_2}, P_{i_3})$, $1 \leq i_1 < i_2 < i_3 \leq r$, are homogeneous, and
3. all triples (p, q, i) , where $i \in \{1, \dots, r\}$ and $p, q \in (P_1 \cup \dots \cup P_r) \setminus P_i$, are good.

Proof. We can assume that $N > (tc_2)^{30d}$, since otherwise the statement is trivial. Since E is semi-algebraic with complexity t , there are polynomials f_1, \dots, f_t of degree at most t , and a Boolean function Φ such that

$$(p_1, p_2, p_3) \in E \iff \Phi(f_1(p_1, p_2, p_3) \geq 0, \dots, f_t(p_1, p_2, p_3) \geq 0) = 1.$$

For each $p, q \in P$ and $i \in \{1, \dots, t\}$, we define the surface $Z_{p,q,i} = \{x \in \mathbb{R}^d : f_i(p, q, x) = 0\}$. Then we set

$$\Sigma = \{Z_{p,q,i} : p, q \in P, 1 \leq i \leq t\}.$$

Thus we have $|\Sigma| = N^2t$. Next we apply Lemma 10 to P and Σ with parameter $\ell = \sqrt{N}$, and obtain subsets Q_1, \dots, Q_ℓ and cells $\Delta_1, \dots, \Delta_\ell$, such that $Q_i \subset \Delta_i$, $|Q_i| = \lfloor \sqrt{N}/4 \rfloor$, and each surface in Σ crosses at most $c_2N^{1/2-1/(4d)}$ cells Δ_i . We note that $c_2 = c_2(d, t)$ is defined in Lemma 10 and $\sqrt{N} \geq \log(tN^2)$. Set $Q = Q_1 \cup \dots \cup Q_\ell$. Each pair $(p, q) \in \binom{Q}{2}$ gives rise to $2t$ surfaces in Σ . By Lemma 10, these $2t$ surfaces cross in total at most $2tc_2N^{1/2-1/(4d)}$ cells Δ_i . Hence there are at most $2tc_2N^{5/2-1/(4d)}$ bad triples of the form (p, q, i) , where $i \in \{1, \dots, \sqrt{N}\}$ and $p, q \in Q \setminus Q_i$. Moreover, there are at most $2tc_2N^{2-1/(4d)}$ bad triples (p, q, i) , where both p and q lie in the same part Q_j and $j \neq i$.

We uniformly at random pick $r = \frac{N^{1/(30d)}}{tc_2}$ subsets (parts) from the collection $\{Q_1, \dots, Q_\ell\}$, and r vertices from each of the subsets that were picked. For a bad triple (p, q, i) with p and q in distinct subsets, the probability that (p, q, i) survives is at most

$$\left(\frac{r}{\sqrt{N}}\right)^3 \left(\frac{r}{\sqrt{N}/4}\right)^2 = \frac{16}{(tc_2)^5} N^{1/(6d)-5/2}.$$

For a bad triple (p, q, i) with p, q in the same subset Q_j , where $j \neq i$, the probability that the triple (p, q, i) survives is at most

$$\left(\frac{r}{\sqrt{N}}\right)^2 \left(\frac{r}{\sqrt{N}/4}\right)^2 = \frac{16}{(tc_2)^4} N^{2/(15d)-2}.$$

Therefore, the expected number of bad triples in our random subset is at most

$$\left(\frac{16}{(tc_2)^5} N^{1/(6d)-5/2}\right) (tc_2N^{5/2-1/(4d)}) + \left(\frac{16}{(tc_2)^4} N^{2/(15d)-2}\right) (tc_2N^{2-1/(4d)}) < 1.$$

Hence we can find disjoint subsets P_1, \dots, P_r , such that $|P_i| \geq r = \frac{N^{1/(30d)}}{tc_2}$, and there are no bad triples (p, q, i) , where $i \in \{1, \dots, r\}$ and $p, q \in (P_1 \cup \dots \cup P_r) \setminus P_i$.

It remains to show that every triple $(P_{i_1}, P_{i_2}, P_{i_3})$ is homogeneous for $1 \leq i_1 < i_2 < i_3 \leq r$. Let $p_1 \in P_{i_1}, p_2 \in P_{i_2}, p_3 \in P_{i_3}$ and suppose $(p_1, p_2, p_3) \in E$. Then for any choice $q_1 \in P_{i_1}, q_2 \in P_{i_2}, q_3 \in P_{i_3}$, we also have $(q_1, q_2, q_3) \in E$. Indeed, since the triple (p_1, p_2, i_3) is good, this implies that $(p_1, p_2, q_3) \in E$. Since the triple (p_1, q_3, i_2) is also good, we have $(p_1, q_2, q_3) \in E$. Finally since (q_2, q_3, i_1) is good, we have $(q_1, q_2, q_3) \in E$. Likewise, if $(p_1, p_2, p_3) \notin E$, then $(q_1, q_2, q_3) \notin E$ for any $q_1 \in P_{i_1}, q_2 \in P_{i_2}, q_3 \in P_{i_3}$. ◀

We are finally ready to prove Theorem 4, which follows immediately from the following theorem.

► **Theorem 15.** *Let P be a set of N points in \mathbb{R}^d and let $E \subset \binom{P}{3}$ be a (symmetric) semi-algebraic relation on P such that E has complexity at most t . If (P, E) is $K_s^{(3)}$ -free, then there exists a subset $P' \subset P$ such that $\binom{P'}{3} \cap E = \emptyset$ and*

$$|P'| \geq 2^{\frac{(\log \log N)^2}{c^s \log \log \log N}},$$

where $c = c(d, t)$.

Proof. The proof is by induction on N and s . The base cases are $s = 3$ or $N \leq (tc_2)^{30d}$, where c_2 is defined in Lemma 10. When $N \leq (tc_2)^{30d}$, the statement holds trivially for sufficiently large $c = c(d, t)$. If $s = 3$, then again the statement follows immediately by taking $P' = P$.

Now assume that the statement holds if $s' \leq s, N' \leq N$ and not both inequalities are equalities. We apply Lemma 14 to (P, E) and obtain disjoint subsets P_1, \dots, P_r , where $r = \frac{N^{1/(30d)}}{tc_2}$, such that $|P_i| \geq \frac{N^{1/(30d)}}{tc_2}$, every triple of parts $(P_{i_1}, P_{i_2}, P_{i_3})$ is homogeneous, and every triple (p, q, i) is good where $i \in \{1, \dots, r\}$ and $p, q \in (P_1 \cup \dots \cup P_r) \setminus P_i$.

Let P_0 be the set of $\frac{N^{1/(30d)}}{tc_2}$ points obtained by selecting one point from each P_i . Since (P_0, E) is $K_s^{(3)}$ -free, we can apply the induction hypothesis on P_0 , and find a set of indices $I = \{i_1, \dots, i_m\}$ such that

$$\log |I| \geq \frac{\left(\log \log \frac{N^{1/(30d)}}{tc_2}\right)^2}{c^s \log \log \log \frac{N^{1/(30d)}}{tc_2}} \geq (1/2) \log \log N,$$

and for every triple $i_1 < i_2 < i_3$ in I all triples with one point in each P_{i_j} does not satisfy E . Hence we have $m = \sqrt{\log N}$, and let $Q_j = P_{i_j}$ for $1 \leq j \leq m$.

For each subset Q_i , we define binary semi-algebraic relations $E_{i,j} \subset \binom{Q_i}{2}$, where $j \neq i$, as follows. Since $E \subset \binom{P}{3}$ is semi-algebraic with complexity t , there are t polynomials f_1, \dots, f_t of degree at most t , and a Boolean function Φ such that $(p_1, p_2, p_3) \in E$ if and only if

$$\Phi(f_1(p_1, p_2, p_3) \geq 0, \dots, f_t(p_1, p_2, p_3) \geq 0) = 1.$$

Fix a point $q_0 \in Q_j$, where $j \neq i$. Then for $p_1, p_2 \in Q_i$, we have $(p_1, p_2) \in E_{i,j}$ if and only if

$$\Phi(f_1(p_1, p_2, q_0) \geq 0, \dots, f_t(p_1, p_2, q_0) \geq 0) = 1.$$

Suppose there are $2^{(\log N)^{1/4}}$ vertices in Q_i that induces a clique in the graph $G_{i,j} = (Q_i, E_{i,j})$. Then these vertices would induce a $K_{s-1}^{(3)}$ -free subset in the original (hypergraph) (P, E) . By the induction hypothesis, we can find a subset $Q'_i \subset Q_i$ such that

$$|Q'_i| \geq 2^{\frac{((1/4) \log \log N)^2}{c^{s-1} \log \log \log N}} \geq 2^{\frac{(\log \log N)^2}{c^s \log \log \log N}},$$

for sufficiently large c , such that $\binom{Q'_i}{3} \cap E = \emptyset$ and we are done. Hence we can assume that each graph $G_{i,j} = (Q_i, E_{i,j})$ has clique number at most $2^{(\log n)^{1/4}}$. By applying Lemma 13 to each Q_i , where Q_i is equipped with $m - 1$ semi-algebraic relations $E_{i,j}$, $j \neq i$, we can find subsets $T_i \subset Q_i$ such that

$$|T_i| \geq \frac{|Q_i|^{1/(c_3 \log m)}}{2^{(\log N)^{1/4} \sqrt{\log N}}} = \frac{2^{\frac{\log N}{30dc_3 \log(\sqrt{\log N})}}}{2^{(\log N)^{3/4}}} \geq 2^{\frac{\log N}{c_5 \log \log N}},$$

where $c_5 = c_5(d, t)$, and $\binom{T_i}{2} \cap E_j = \emptyset$ for all $j \neq i$. Therefore, we now have subsets T_1, \dots, T_m , such that

1. $m = \sqrt{\log N}$,
2. for any triple $(T_{i_1}, T_{i_2}, T_{i_3})$, $1 \leq i_1 < i_2 < i_3 \leq m$, every triple with one vertex in each T_{i_j} is not in E ,
3. for any pair (T_{i_1}, T_{i_2}) , $1 \leq i_1 < i_2 \leq m$, every triple with two vertices T_{i_1} and one vertex in T_{i_2} is not in E , and every triple with two vertices T_{i_2} and one vertex in T_{i_1} is also not in E .

By applying the induction hypothesis to each (T_i, E) , we obtain a collection of subsets $U_i \subset T_i$ such that

$$\log |U_i| \geq \frac{\left(\log\left(\frac{\log N}{c_5 \log \log N}\right)\right)^2}{c^s \log \log\left(\frac{\log N}{c_5 \log \log N}\right)} \geq \frac{(\log \log N - \log(c_5 \log \log N))^2}{c^s \log \log \log N},$$

and $\left(\bigcup_{i=1}^m U_i\right) \cap E = \emptyset$. Let $P' = \bigcup_{i=1}^m U_i$. Then by above we have $\binom{P'}{3} \cap E = \emptyset$ and

$$\begin{aligned} \log |P'| &\geq \frac{(\log \log N - \log(c_5 \log \log N))^2}{c^s \log \log \log N} + \frac{1}{2} \log \log N \\ &\geq \frac{(\log \log N)^2 - 2(\log \log N) \log(c_5 \log \log N) + (\log(c_5 \log \log N))^2}{c^s \log \log \log N} + \frac{1}{2} \log \log N \\ &\geq \frac{(\log \log N)^2}{c^s \log \log \log N}, \end{aligned}$$

for sufficiently large $c = c(d, t)$. ◀

5 Application: One-sided hyperplanes

Let us consider a finite set H of hyperplanes in \mathbb{R}^d in general position, that is, every d members in H intersect at a distinct point. Let $OSH_d(s, n)$ denote the smallest integer N such that every set H of N hyperplanes in \mathbb{R}^d in general position contains s members H_1 such that the vertex set of the arrangement of H_1 lies above the $x_d = 0$ hyperplane, or contains n members H_2 such that the vertex set of the arrangement of H_2 lies below the $x_d = 0$ hyperplane.

In 1992, Matoušek and Welzl [28] observed that $OSH_2(s, n) = (s-1)(n-1)+1$. Dujmović and Langerman [15] used the existence of $OSH_d(n, n)$ to prove a ham-sandwich cut theorem for hyperplanes. Again by adapting the Erdős-Rado argument, Conlon et al. [13] showed that for $d \geq 3$,

$$OSH_d(s, n) \leq \text{twr}_{d-1}(c_6 s n \log n), \tag{5}$$

where c_6 is a constant that depends only on d . See Eliáš and Matoušek [17] for more related results, including lower bound constructions.

Since each hyperplane $h_i \in H$ is specified by the linear equation

$$a_{i,1}x_1 + \dots + a_{i,d}x_d = b_i,$$

we can represent $h_i \in H$ by the point $h_i^* \in \mathbb{R}^{d+1}$ where $h_i^* = (a_{i,1}, \dots, a_{i,d}, b_i)$ and let $P = \{h_i^* : h_i \in H\}$. Then we define a relation $E \subset \binom{P}{d}$ such that $(h_{i_1}^*, \dots, h_{i_d}^*) \in E$ if and only if $h_{i_1} \cap \dots \cap h_{i_d}$ lies above the hyperplane $x_d = 0$ (i.e. the d -th coordinate of the intersection point is positive). Clearly, E is a semi-algebraic relation with complexity at most $t = t(d)$. Therefore, as an application of Theorem 4 and Corollary 5, we make the following improvement on (5).

► **Theorem 16.** For fixed $s \geq 4$, we have $OSH_3(s, n) \leq 2^{n^{o(1)}}$. For fixed $d \geq 4$ and $s \geq d+1$, we have

$$OSH_d(s, n) \leq \text{twr}_{d-1}(n^{o(1)}).$$

6 Monochromatic triangles

In this section, we will prove Theorem 6.

Proof of Theorem 6. We proceed by induction on m . The base case when $m = 1$ is trivial. Now assume that the statement holds for $m' < m$. Set $N = 2^{cm \log \log m}$, where $c = c(d, t)$ will be determined later, and let $E_1, \dots, E_m \subset \binom{P}{2}$ be (symmetric) semi-algebraic relations on P such that $\binom{P}{2} = E_1 \cup \dots \cup E_m$, and each E_i has complexity at most t . For sake of contradiction, suppose P does not contain three points such that every pair of them is in E_i for some fixed i .

For each relation E_i , there are t polynomials $f_{i,1}, \dots, f_{i,t}$ of degree at most t , and a Boolean function Φ_i such that

$$(p, q) \in E_i \quad \Leftrightarrow \quad \Phi_i(f_{i,1}(p, q) \geq 0, \dots, f_{i,t}(p, q) \geq 0) = 1.$$

For $1 \leq i \leq m, 1 \leq j \leq t, p \in P$, we define the surface $Z_{i,j,p} = \{x \in \mathbb{R}^d : f_{i,j}(p, x) = 0\}$, and let

$$\Sigma = \{Z_{i,j,p} : 1 \leq i \leq m, 1 \leq j \leq t, p \in P\}.$$

Hence $|\Sigma| = mtN$. We apply Lemma 9 to Σ with parameter $r = 2tm$, and decompose \mathbb{R}^d into $c_1(2tm)^{2d}$ regions Δ_i , where $c_1 = c_1(t, d)$ is defined in Lemma 9, such that each region Δ_i is crossed by at most $tmN/r = N/2$ members in Σ . By the pigeonhole principle, there is a region $\Delta \subset \mathbb{R}^d$, such that $|\Delta \cap P| \geq \frac{N}{c_1(2tm)^{2d}}$, and at most $N/2$ members in Σ crosses Δ . Let P_1 be a set of exactly $\lfloor \frac{N}{c_1(2tm)^{2d}} \rfloor$ points in $P \cap \Delta$, and let P_2 be the set of points in $P \setminus P_1$ that does not give rise to a surface that crosses Δ . Hence

$$|P_2| \geq N - \frac{N}{c_1(2tm)^{2d}} - \frac{N}{2} \geq \frac{N}{4}.$$

Therefore, each point $p \in P_2$ has the property that $p \times P_1 \subset E_i$ for some fixed i . We define the function $\chi : P_2 \rightarrow \{1, \dots, m\}$, such that $\chi(p) = i$ if and only if $p \times P_1 \subset E_i$. Set $I = \{\chi(p) : p \in P_2\}$ and $m_0 = |I|$, that is, m_0 is the number of distinct relations (colors) between the sets P_1 and P_2 . Now the proof falls into 2 cases.

Case 1. Suppose $m_0 > \log m$. By the assumption, every pair of points in P_1 is in E_i for some $i \in \{1, \dots, m\} \setminus I$. By the induction hypothesis, we have

$$\frac{2^{cm \log \log m}}{c_1(2tm)^{2d}} \leq |P_1| \leq 2^{c(m-m_0) \log \log m}.$$

Hence

$$cm_0 \log \log m \leq \log(c_1(2tm)^{2d}) \leq 2d \log(c_1 2tm),$$

which implies

$$m_0 \leq \frac{2d \log(c_1 2tm)}{c \log \log m},$$

and we have a contradiction for sufficiently large $c = c(d, t)$.

Case 2. Suppose $m_0 \leq \log m$. By the pigeonhole principle, there is a subset $P_3 \subset P_2$, such that $|P_3| \geq \frac{N}{4m_0}$ and $P_1 \times P_3 \subset E_i$ for some fixed i . Hence every pair of points $p, q \in P_3$ satisfies $(p, q) \notin E_i$, for some fixed i . By the induction hypothesis, we have

$$\frac{2^{cm \log \log m}}{4m_0} \leq |P_3| \leq 2^{c(m-1) \log \log m}.$$

Therefore

$$c \log \log m \leq \log(4m_0) \leq \log(4 \log(m)),$$

which is a contradiction since c is sufficiently large. This completes the proof of Theorem 6. ◀

6.1 Lower bound construction and Schur numbers

Before we prove Theorem 7, let us recall a classic Theorem of Schur [31] which is considered to be one of the earliest applications of Ramsey Theory. A subset of numbers $P \subset \mathbb{R}$ is said to be *sum-free* if for any two (not necessarily distinct) elements $x, y \in P$, their sum $x + y$ is not in P . The Schur number $S(m)$ is defined to be the maximum integer N for which the integers $\{1, \dots, N\}$ can be partitioned into m sum-free sets.

Given a partition $\{1, \dots, N\} = P_1 \cup \dots \cup P_m$ into m parts such that P_i is sum-free, we can define an m -coloring on the edges on a complete $(N + 1)$ -vertex graph which does not contain a monochromatic triangle as follows. Let $V = \{1, \dots, N + 1\}$ be the vertex set, and we define the coloring $\chi : \binom{V}{2} \rightarrow m$ by $\chi(x, y) = i$ iff $|x - y| \in P_i$. Now suppose for sake of contradiction there are vertices x, y, z that induces a monochromatic triangle, say with color i , such that $x < y < z$. Then we have $y - x, z - y, z - x \in P_i$ and $(y - x) + (z - y) = (z - x)$, which is a contradiction since P_i is sum free. Therefore $S(m) < R_2(3; m)$.

Since Schur’s original 1916 paper, the lower bound on $S(m)$ has been improved by several authors [2, 1, 22], and the current record of $S(m) \geq \Omega(3.19^m)$ is due to Fredricksen and Sweet [24]. Their lower bound follows by computing $S(6) \geq 538$, and using the recursive formula

$$S(m) \geq c_\ell(2S(\ell) + 1)^{m/\ell},$$

which was established by Abbott and Hanson [1]. Fredricksen and Sweet also computed $S(7) \geq 1680$, which we will use to prove Theorem 7.

► **Lemma 17.** *For each integer $\ell \geq 1$, there is a set P_ℓ of $(1681)^\ell$ points in \mathbb{R} equipped with semi-algebraic relations $E_1, \dots, E_{7\ell} \subset \binom{P_\ell}{2}$, such that*

1. $E_1 \cup \dots \cup E_{7\ell} = \binom{P_\ell}{2}$,
2. E_i has complexity at most 5000,
3. E_i is translation invariant, that is, $(x, y) \in E_i$ iff $(x + C, y + C) \in E_i$, and
4. the graph $G_{\ell,i} = (P_\ell, E_i)$ is triangle free for all i .

Proof. We start by setting $P_1 = \{1, 2, \dots, 1681\}$. By [24], there is a partition on $\{1, \dots, 1680\} = A_1 \cup \dots \cup A_7$ into seven parts, such that each A_i is sum-free. For $i \in \{1, \dots, 7\}$, we define the binary relation E_i on P_1 by

$$(x, y) \in E_i \iff (1 \leq |x - y| \leq 1680) \wedge (|x - y| \in A_i).$$

Since $|A_i| \leq 1680$, E_i has complexity at most 5000. By the arguments above, the graph $G_{1,i} = (P_1, E_i)$ is triangle free for all $i \in \{1, \dots, 7\}$. In what follows, we blow-up this construction so that the statement holds.

Having defined $P_{\ell-1}$ and $E_1, \dots, E_{7\ell-7}$, we define P_ℓ and $E_{\ell-6}, \dots, E_\ell$ as follows. Let $C = C(\ell)$ be a very large constant, say $C > (5000 \cdot \max\{P_{\ell-1}\})^2$. We construct 1681

translated copies of $P_{\ell-1}$, $Q_i = P_{\ell-1} + iC$ for $1 \leq i \leq 1681$, and set $P_\ell = Q_1 \cup \dots \cup Q_{1681}$. For $1 \leq j \leq 7$, we define the relation $E_{\ell-7+j}$ by

$$(x, y) \in E_{\ell-7+j} \quad \Leftrightarrow \quad (C/2 \leq |x - y| \leq 1682C) \wedge (\exists z \in A_j : ||x - y|/C - z| < 1/1000).$$

Clearly $E_1, \dots, E_{7\ell}$ satisfy properties (1), (2), and (3). The fact that $G_{\ell,i} = (P_\ell, E_i)$ is triangle follows from the same argument as above. \blacktriangleleft

Theorem 7 immediately follows from Lemma 17.

References

- 1 H. L. Abbott and D. Hanson. A problem of schur and its generalizations. *Acta Arith.*, 20:175–187, 1972.
- 2 H. L. Abbott and L. Moser. Sum-free sets of integers. *Acta Arith.*, 11:392–396, 1966.
- 3 P. K. Agarwal and J. Erickson. Optimal partition trees. In *In Proc. 26th Ann. ACM Sympos. Comput. Geom.*, pages 1–10, 2010.
- 4 P. K. Agarwal and J. Erickson. Geometric range searching and its relatives. In J. E. Goodman B. Chazelle and R. Pollack, editors, *Advances in Discrete and Computational Geometry*, pages 1–56, 1998.
- 5 M. Ajtai, J. Komlós, and E. Szemerédi. A note on ramsey numbers. *J. Combin. Theory Ser. A*, 29:354–360, 1980.
- 6 N. Alon, J. Pach, R. Pinchasi, R. Radoičić, and M. Sharir. Crossing patterns of semi-algebraic sets. *J. Combin. Theory Ser. A*, 111:310–326, 2005.
- 7 S. Basu, R. Pollack, and M. F. Roy. *Algorithms in Real Algebraic Geometry*. Springer-Verlag, Berlin, 2nd edition edition, 2006.
- 8 T. Bohman. The triangle-free process. *Adv. Math.*, 221:1653–1677, 2009.
- 9 T. Bohman and P. Keevash. The early evolution of the h -free process. *Invent. Math.*, 181:291–336, 2010.
- 10 B. Bukh and M. Matoušek. Erdős-Szekeres-type statements: Ramsey function and decidability in dimension 1. *Duke Math. Journal*, 63:2243–2270, 2014.
- 11 B. Chazelle, H. Edelsbrunner, L. Guibas, and M. Sharir. A singly exponential stratification scheme for real semi-algebraic varieties and its applications. *Theor. Comput. Sci.*, 84:77–105, 1991.
- 12 K. L. Clarkson and P. W. Shor. Applications of random sampling in computational geometry, ii. *Discrete Comput. Geom.*, 4:387–421, 1989.
- 13 D. Conlon, J. Fox, J. Pach, B. Sudakov, and A. Suk. Ramsey-type results for semi-algebraic relations. *Trans. Amer. Math. Soc.*, 366:5043–5065, 2014.
- 14 D. Conlon, J. Fox, and B. Sudakov. Hypergraph ramsey numbers. *J. Amer. Math. Soc.*, 23:247–266, 2010.
- 15 V. Dujmović and S. Langerman. A center transversal theorem for hyperplanes and applications to graph drawing. In *In Proc. 27th Ann. ACM Sympos. Comput. Geom.*, pages 117–124, 2011.
- 16 M. Eliáš, J. Matoušek, E. Roldán-Pensado, and Z. Safernová. Lower bounds on geometric Ramsey functions. *SIAM J. Discrete Math*, 28:1960–1970, 2014.
- 17 M. Eliáš and J. Matoušek. Higher-order Erdős-Szekeres theorems. *Advances in Mathematics*, 244:1–15, 2013.
- 18 P. Erdős. Some remarks on the theory of graphs. *Bull. Amer. Math. Soc.*, 53:292–294, 1947.
- 19 P. Erdős, A. Hajnal, and R. Rado. Partition relations for cardinal numbers. *Acta Math. Acad. Sci. Hungar.*, 16:93–196, 1965.

- 20 P. Erdős and R. Rado. Combinatorial theorems on classifications of subsets of a given set. *Proc. London Math. Soc.*, 3:417–439, 1952.
- 21 P. Erdős and G. Szekeres. A combinatorial problem in geometry. *Compos. Math.*, 2:463–470, 1935.
- 22 G. Exoo. A Lower Bound for Schur numbers and multicolor Ramsey numbers of K_3 . *Electronic J. Combinatorics*, 1:1–3, 1994.
- 23 J. Fox, J. Pach, B. Sudakov, and A. Suk. Erdős-Szekeres-type theorems for monotone paths and convex bodies. *Proceedings of the London Mathematical Society*, 105:953–982, 2012.
- 24 H. Fredricksen and M. Sweet. Symmetric sum-free partitions and lower bounds for schur numbers. *Electronic J. Combinatorics*, 7:1–9, 2000.
- 25 J. H. Kim. The ramsey number $r(3, t)$ has order of magnitude $t^2/\log t$. *Random Structures Algorithms*, 7:173–207, 1995.
- 26 V. Koltun. Almost tight upper bounds for vertical decompositions in four dimensions. *J. ACM*, 51:699–730, 2004.
- 27 J. Matoušek. Efficient partition trees. *Discrete Comput. Geom.*, 8:315–334, 1992.
- 28 J. Matoušek and E. Welzl. Good splitters for counting points in triangles. *J. Algorithms*, 13:307–319, 1992.
- 29 J. Milnor. On the betti numbers of real varieties. *Proc. Amer. Math. Soc.*, 15:275–280, 1964.
- 30 D. Mubayi and A. Suk. A ramsey-type result for geometric ℓ -hypergraphs. *European Journal of Combinatorics*, 41:232–241, 2014.
- 31 I. Schur. Über die Kongruenz $x^m + y^m = z^m \pmod p$. *Jahresber. Deutch. Math. Verein.*, 25:114–117, 1916.
- 32 M. J. Steele. Variations on the monotone subsequence theme of Erdős and Szekeres. In D. Aldous, editor, *Discrete Probability and Algorithms, IMA Volumes in Mathematics and its Applications*, pages 111–131, Berlin, 1995. Springer.
- 33 A. Suk. A note on order-type homogeneous point sets. *Mathematika*, 60:37–42, 2014.
- 34 R. Thom. Sur l’homologie des variétés algébriques réelles. In *Differential and Combinatorial Topology (A Symposium in Honor of Marston Morse)*, pages 255–265, Princeton, N.J., 1965. Princeton University.
- 35 H. E. Warren. Lower bounds for approximation by nonlinear manifold. *Trans. Amer. Math. Soc.*, 133:167–178, 1968.